

Weight Management Over the Reproductive Years for Adult Women Living with Obesity

Helena Piccinini-Vallis MD PhDⁱ, Kristi Adamo PhDⁱⁱ,
Rhonda Bell PhDⁱⁱⁱ, Leticia Pereira PhD^{iv}, Kara Nerenberg MD^v

- i) Department of Family Medicine, Dalhousie University
- ii) Faculty of Health Science, University of Ottawa
- iii) Division of Human Nutrition, University of Alberta
- iv) Department of Agricultural, Food and Nutritional Sciences, University of Alberta
- v) Department of Medicine, University of Calgary; Foothills Medical Centre

Cite this Chapter

Piccinini-Vallis H., Adamo K, Bell R, Pereira L, Nerenberg K. Canadian Adult Obesity Clinical Practice Guidelines: Weight Management Over the Reproductive Years for Adult Women Living with Obesity. Available from: <https://obesitycanada.ca/guidelines/reproductive>. Accessed [date].

Update History

Version 1, August 4, 2020. The Canadian Adult Obesity Clinical Practice Guidelines are a living document, with only the latest chapters posted at obesitycanada.ca/guidelines.

KEY MESSAGES FOR PRIMARY CARE PROVIDERS

This chapter addresses the management of weight related to three phases of a woman's reproductive years – preconception, during pregnancy and postpartum – for adult women with obesity. Although these reproductive periods are addressed separately, it is important to consider that these phases represent the continuum of weight management over

the reproductive years in women with obesity. During these time periods, women frequently access the healthcare system, thus providing clinicians with health promotion opportunities which may have positive impacts on the short- and long-term health of both the woman and her children. Discussion of the obstetric and anesthetic management for women with obesity during pregnancy is beyond the scope of this clinical practice guideline.

RECOMMENDATIONS

These recommendations pertain to the management of weight over the reproductive years for adult women with obesity (i.e., body mass index [BMI] ≥ 30 kg/m²) with a singleton pregnancy, who are ≥ 18 years of age and do not have pre-existing diabetes or gestational diabetes.

1. General advice: We recommend primary care providers should discuss weight management targets specific to the

reproductive years with adult women with obesity: preconception weight loss (Level 3, Grade C),¹⁻⁴ gestational weight gain of 5 kg–9 kg over the entire pregnancy (Level 4, Grade D);⁵ postpartum weight loss of – at minimum – gestational weight gain (Level 3, Grade C)^{6,7} to reduce the risk of adverse outcomes in the current or in a future pregnancy.

2. Combined behaviour change interventions: Primary care providers should offer behaviour change interventions, including both nutrition and physical activity, to adult

women with obesity who are considering a pregnancy (Level 3, Grade C),^{7,8} who are pregnant (Level 2a, Grade B)⁹⁻¹⁵ and who are postpartum (Level 1a, Grade A)¹⁶ in order to achieve weight targets.

3. Nutrition counselling alone: We recommend primary care providers encourage and support pregnant women with obesity to consume foods consistent with a healthy dietary pattern in order to meet their target gestational weight gain (Level 3, Grade C).¹⁷

4. Physical activity counselling alone: We recommend primary care providers encourage and support pregnant women with obesity who do not have contraindications to exercise

during pregnancy to engage in at least 150 minutes per week of moderate intensity physical activity to assist in the management of gestational weight gain (Level 3, Grade C).¹⁸⁻²¹

5. Pharmacotherapy: Healthcare providers should not prescribe metformin for managing gestational weight gain in women with obesity (Level 1b, Grade A).²²⁻²⁴ We suggest no weight management medications during pregnancy or breastfeeding (Level 4, Grade D).²⁵

6. Breastfeeding: We recommend women with obesity be offered additional breastfeeding support due to decreased rates of initiation and continuation (Level 3, Grade C).²⁶

KEY MESSAGES FOR WOMEN WITH OBESITY DURING THE REPRODUCTIVE YEARS

The reproductive years, including before, during and after pregnancy, bring many additional challenges for women with obesity in maintaining a healthy weight beyond eating well and being physically active. It is important for women with obesity to seek advice and support from their healthcare providers on strategies to optimize their own health outcomes, as well as those of their children, over both the short and long-term.

The strategies described in this chapter include:

- 1) Entering pregnancy at a lower BMI;
- 2) Targeting weight gain during the entire pregnancy to 5 kg–9 kg; and
- 3) Returning to at least the pre-pregnancy BMI in the year after delivery.

Preconception period

Background

A woman's preconception health status, particularly the control of pre-pregnancy medical co-morbidities, has been demonstrated to markedly impact the maternal, obstetrical and fetal health outcomes during pregnancy.^{27,28} While many pregnancies are unplanned, a preconception consultation offers women considering a pregnancy the opportunity to engage in discussions with their healthcare providers about their individualized health risks during pregnancy, and to make informed decisions. In addition, the preconception period is a time when interventions can be implemented to reduce the risks of common adverse pregnancy outcomes.^{27,28}

As outlined in Table 1, for women with obesity who become pregnant, a BMI of ≥ 30 kg/m² has been consistently associated with the following: infertility,^{29,30} congenital malformations,³¹ and numerous clinically important adverse pregnancy outcomes (in a dose-response relationship by BMI) including hypertensive disorders of pregnancy, gestational diabetes, Caesarean delivery, preterm birth, etc.^{2,31-33} While there remains a paucity of data to specifically guide clinicians on how best to counsel women with

obesity about these pregnancy-associated health risks, emerging data demonstrate that weight management prior to pregnancy may reduce many of the above risks.¹ Discussion of specific nutritional supplementation (e.g., folic acid) and obstetrical care is beyond the scope of this clinical practice guideline.

Clinical strategies and resources to achieve preconception weight management for women with obesity

Combined behavioural interventions: There are few trials specifically conducted in women with obesity planning a pregnancy. Agha et al.'s systematic review⁸ of randomized controlled trials (RCTs) in preconception women with overweight or obesity found that combined behavioural health interventions (i.e., nutrition and physical activity) were associated with significant reductions in gestational weight gain once pregnant. This is similar to the findings of Krukowski et al.⁷ who examined the effectiveness of a combined behavioural health intervention (combined nutrition and physical activity) beginning in the preconception period or up to 10 weeks' gestational age. The study found that women with obesity in the intervention arm had reduced gestational weight gain at 36 weeks' gestation. In addition, as outlined in the postpartum period section, the use of combined behaviour change interventions in the postpartum period was associated with greater

Table 1: Associations between obesity, gestational weight gain and adverse clinical outcomes over the reproductive (preconception, antenatal, and postpartum) periods

Reproductive Period	Weight Management Issue	Adverse Clinical Outcome
Preconception	Pre-pregnancy obesity	<ul style="list-style-type: none"> ↑ Infertility²⁹ ↑ Gestational diabetes³⁴ ↑ Hypertensive disorders of pregnancy (i.e., gestational hypertension and preeclampsia)^{2,4,32,35,36} ↑ Maternal venous thromboembolism^{32,37} ↑ Postpartum depression³ ↑ Miscarriage²⁸ ↑ Caesarean delivery³² ↑ Congenital malformations^{31,38} ↑ Newborn asphyxia³³ ↑ Macrosomia/large-for-gestational-age³³
Pregnancy	Excess gestational weight gain	<ul style="list-style-type: none"> ↑ Gestational diabetes^{38,39} ↑ Hypertensive disorders of pregnancy (i.e., Gestational hypertension and preeclampsia)^{36,40–42} ↑ Caesarean delivery^{1,40,42,43} ↑ Preterm birth (medically indicated)⁴³ ↑ Macrosomia^{40,42,44,45} ↑ Neonatal hyperinsulinemia⁴⁵ ↑ Neonatal hypoglycemia, hypomagnesemia, & hypocalcemia⁴⁴ ↑ Postpartum weight retention^{46–49}
	Low gestational weight gain or weight loss	<ul style="list-style-type: none"> ↓ Hypertensive disorders of pregnancy (i.e., Gestational Hypertension³⁸ and Preeclampsia³⁵) ↓ Caesarean delivery^{1,40,43,50} ↓ Large for gestational age infant^{42,43,51,52} ↑ Macrosomia^{2,50}
	Weight loss	<ul style="list-style-type: none"> ↑ Low birth weight infant⁴⁰
Postpartum	Pre-pregnancy obesity	<ul style="list-style-type: none"> ↑ Postpartum weight retention³²
	Obesity during pregnancy	<ul style="list-style-type: none"> ↑ Obesity by 9 months postpartum⁵³
	Excess postpartum weight retention	<ul style="list-style-type: none"> ↑ Gestational diabetes in future pregnancy¹ ↑ Hypertensive disorders in a future pregnancy¹
	Reduction in BMI by 2 kg/m ²	<ul style="list-style-type: none"> ↓ Gestational diabetes in future pregnancy⁶

postpartum weight loss, which may impact the health outcomes of future pregnancies.

Pharmacotherapy: At present, the fetal effects of several pharmacologic agents used for the purpose of weight management are not known. Therefore, in general, it is not recommended that these agents be used during pregnancy. Consideration should be given to stopping these medications prior to pregnancy to limit exposure to the developing fetus.²⁵

Pregnancy/antenatal period

Background

Women with obesity have an increased risk of numerous adverse maternal, obstetrical and fetal outcomes during pregnancy, as outlined in Table 1. One promising strategy to reduce or prevent these adverse outcomes is through the achievement of the recommended guidelines for gestational weight gain during pregnancy (see below), though this remains an area of active study.

The current recommended guidelines for weight gain during pregnancy for uncomplicated singleton pregnancies were published by the Institute of Medicine (IOM) in 2009⁵⁴ and adopted by Health Canada in 2010⁵⁴ (Table 2). These recommendations were developed based on the findings of numerous observational studies. These studies consistently demonstrate that gestational weight gain above or below these recommended ranges is associated with several important adverse clinical outcomes for women

and their offspring including: birthweight, large for gestational age, small for gestational age, Caesarean delivery, preterm delivery, postpartum maternal weight retention, and childhood obesity.⁵¹ However, the data on the association between gestational weight gain and small-for-gestational-age infants remain contradictory.^{40,43,50,51}

Several factors influence gestational weight gain for women with obesity, including depression,⁵⁵ health behaviours,⁵⁶ patient expectations and knowledge,⁵⁷ educational attainment/socioeconomic status,⁵⁸ maternal age, parity and ethnicity.⁵⁹ Importantly, obesity prior to pregnancy is also an independent risk factor for excess gestational weight gain.^{60–63} Although there are no data specifically for women with obesity, there is evidence that advice from a prenatal healthcare provider is both desired by women⁶⁴ and can positively influence gestational weight gain.⁶⁵

Considerations for weight management during pregnancy for women with obesity

Women with a singleton pregnancy can expect to gain approximately 8.5 kilograms during the course of a full-term pregnancy, regardless of the increase in their own adipose tissue mass, as a result of the following physiologic increases in weight: term baby (approximately 3 kg); the placenta; amniotic fluid; uterine muscle; increase in intravascular blood volume; and the increase in total body water volume. For women with obesity, this weight gain amount is just below the upper limit of the IOM/Health Canada recommendations for optimal gestational weight gain (5kg–9 kg throughout the entire pregnancy). As a result, weight management

Table 2: Gestational weight gain recommendations (Institute of Medicine, 2009)

Pre-pregnancy BMI	Total Weight Gain		Rates of Weight Gain* 2nd and 3rd Trimester	
	Range in kg	Range in lbs	Mean (range) in kg/week	Mean (range) in lbs/week
Underweight (< 18.5 kg/m ²)	12.5–18	28–40	0.51 (0.44–0.58)	1 (1–1.3)
Normal weight (18.5–24.9 kg/m ²)	11.5–16	25–35	0.42 (0.35–0.50)	1 (0.8–1)
Overweight (25.0–29.9 kg/m ²)	7–11.5	15–25	0.28 (0.23–0.33)	0.6 (0.5–0.7)
Obesity (> 30.0 kg/m ²)	5–9	11–20	0.22 (0.17–0.27)	0.5 (0.4–0.6)

Source: Institute of Medicine and National Research Council. 2009. Weight Gain During Pregnancy: Reexamining the Guidelines. <https://doi.org/10.17226/12584>. Reproduced with permission from the National Academy of Sciences, courtesy of the National Academies Press, Washington, D.C.

can be challenging for women with obesity, and should therefore be addressed at the first prenatal appointment and throughout pregnancy. An evidence-based point-of-care tool, [The 5As of Healthy Pregnancy Weight Gain](#), is available to support primary care providers in discussing healthy pregnancy weight gain with women.⁶⁶

Advice from prenatal healthcare providers positively influences women's achievement of gestational weight gain targets.⁶⁷⁻⁷⁰ Further, pregnant women report that they want to discuss gestational weight gain with their healthcare providers.⁷¹⁻⁷³ Although Canadian data show that most healthcare providers reported routinely weighing pregnant women, only a minority routinely discussed the actual weight results.^{74,75} Weight-related discussions are often perceived by healthcare providers as sensitive in nature and are therefore often avoided.⁷⁶⁻⁷⁹ This hesitation in raising the topic could be mitigated to some extent by having a good patient-provider relationship.⁸⁰ Primary care providers have a longitudinal relationship with their patients and are therefore in an advantageous position to be able to support women with obesity to achieve the recommended targets for gestational weight gain. A theoretical framework that is highly relevant to primary care clinicians is the Patient-Centred Clinical Method,⁸¹ comprised of four interactive domains:

- a) Exploring women's experience of changes in weight during pregnancy;
- b) Understanding women's proximal and distal contexts;
- c) Finding common ground on the best approach to support appropriate weight gain during pregnancy; and
- d) Enhancing the patient-clinician relationship.

From a patient-centred perspective, it is key to address the first two domains. Otherwise, the provider's recommendations are often neither meaningful nor able to be translated into patient behaviour change. (See the [Effective Psychological and Behavioural Interventions for People Living with Obesity](#) chapter).

Clinical strategies and resources to achieve recommended gestational weight gain targets for women with obesity

Based upon the above, a number of interventions and models of care could be helpful to women with obesity during pregnancy, when approached from a patient-centred perspective.

Nutrition counselling: When implemented early in pregnancy, nutrition counselling can assist women with obesity in managing their pregnancy weight gain. Wolff et al.¹⁷ randomized pregnant women with obesity to either an intervention arm (n=23), consisting of 10 one-hour sessions with a trained dietitian, or usual care (n=27). The sessions focused on providing advice on eating a healthy diet according to Danish national dietary recommendations, with the use of food records to identify unhealthy eating patterns and to give individualized feedback for improvement. At 36 weeks' gestation, the mean gestational weight gain for the

intervention group was 6.6 kg, compared to a mean gestational weight gain of 13.3 kg for the control group (p=0.002).

Physical activity: Physical activity (a term used synonymously with exercise in this chapter) during pregnancy has been shown to be beneficial to women with obesity in managing their pregnancy weight gain ([2019 Canadian Guideline for Physical Activity Throughout Pregnancy](#)). Daly et al.²⁰ compared total gestational weight gain at 36 weeks' gestation between 88 pregnant women assigned to either an intervention consisting of intensive, medically supervised exercise or to usual care. Fewer women in the exercise intervention group gained weight in excess of the guidelines compared to women in the control group (23.5% versus 45.2%; p<0.05). Barakat et al.¹⁹ compared total gestational weight gain between pregnant women randomized to an exercise intervention or to standard care as an analysis of a secondary outcome in a RCT. The intervention consisted of 50- to 55-minute training sessions (three days per week) from nine to 11 weeks' gestation until 38 to 39 weeks' gestation (85 sessions in total). Among women with obesity (n=54), 44.0% of women in the intervention group experienced excess gestational weight gain compared with 51.7% of women in the control group. In a case-control study undertaken by Claesson et al.,²¹ the intervention consisted of motivational talks to assist with behaviour change and regular exercise (aqua aerobics). The intervention group experienced significantly lower weight gain than the control group (8.7 kg vs 11.3 kg; p<0.001).

Combined behavioural interventions: A comprehensive approach targeting both nutrition and exercise has been shown to be effective in managing gestational weight gain. Vinter et al.¹³ randomized 360 women with obesity at 10 to 14 weeks' gestation to a comprehensive lifestyle intervention or standard care (Lifestyle in Pregnancy Study: LiP). Women in the intervention group received four 30- to 60-minute diet counselling sessions delivered by trained dietitians and were encouraged to be moderately physically active for 30 to 60 minutes daily. At 35 weeks' gestation, women in the intervention group had gained significantly less weight than women in the control group (7.4 kg versus 8.6 kg, p=0.014).

Petrella et al.¹⁴ randomized 38 women with obesity to a therapeutic lifestyle changes program consisting of 1800 kcal/day and 30 minutes of physical activity of moderate intensity on at least three days per week, or to usual care. At delivery, women in the intervention group experienced significantly lower gestational weight gain than women in the control group (6.7 kg versus 10.1 kg; p=0.047). In addition, 77.8% of women receiving the intervention experienced guideline-concordant gestational weight gain, compared with 30.0% of women in the control group (p=0.003).

Poston et al.¹⁰ randomly assigned 1,555 pregnant women with obesity who were between 15- and 18-weeks' gestation to either a theory-based intensive behavioural intervention or to standard antenatal care (UK Pregnancies Better Eating and Activity Trial: UPBEAT). The intervention consisted of eight 60-minute weekly sessions addressing nutrition and physical activity by setting and achieving SMART goals. At delivery (mean gestational age 39.5

weeks for both groups), women in the intervention group had gained less weight than women in the control group (7.19 kg versus 7.76 kg, $p=0.041$).

Haby et al.¹¹ conducted an RCT evaluating a prenatal lifestyle intervention compared to usual care for women with obesity, with gestational weight gain as one of the outcomes. The intervention group ($n=459$) received two 30-minute midwife-delivered personalized counselling sessions on food and physical activity during early pregnancy. Individualized dietary advice from a dietitian, food discussion groups with a dietitian, aqua aerobics led by a physiotherapist, prescriptions for physical activity, walking poles, pedometers and information about community resources were also available to the intervention group if desired. The control group ($n=895$) received usual care. Per-protocol analysis showed significantly lower gestational weight gain in the intervention group compared to the control group (8.9 kg vs 11.2 kg; $p=0.031$).

Renault et al.⁸² undertook an RCT with 425 women with obesity being assigned to one of three groups: physical activity plus diet, physical activity alone, or control. Both interventions resulted in less gestational weight gain than the control ($p = 0.008$). The median gestational weight gain, compared with the control group (10.9 kg), was lower in both intervention groups: physical activity plus diet 8.6 kg ($p=0.01$), and physical activity alone group 9.4 kg ($p=0.042$).

Metformin: To date, RCT evidence for the use of metformin in the management of gestational weight gain for women with obesity who do not have diabetes is conflicting, and therefore metformin is not recommended. Although a Cochrane review by Dodd et al.⁸³ of three studies of metformin (up to 3000 mg/day) in pregnant women with overweight or obesity found that women who received metformin may have a slightly lower gestational weight gain compared with placebo, the heterogeneity of the studies prevented meaningful evaluations.

Healthcare provider behaviour: There are limited data specifically focused on pregnant women with obesity. Observational studies in the general population show that healthcare provider patient-centredness influences patients' adherence to weight gain recommendations,^{84–86} patients' perception of "finding common ground",⁸¹ increased patient satisfaction and decreased burden of symptoms.⁸⁷

In a qualitative Canadian study, Nikolopoulos et al. (2017)⁷⁴ concluded that healthcare providers should initiate discussions about gestational weight gain in a patient-centred manner, specifically by "asking women how they feel about discussing weight".

Lindberg et al.⁸⁸ examined the weight outcomes before and after implementation of a 'best practice alert' in the electronic medical record, which provided tailored gestational weight gain goals and patient education materials based on patients' pre-pregnancy BMI and the 2009 IOM guidelines for weight gain during pregnancy. Overall, the intervention was associated with a significant increase

in the proportion of women with obesity who had guideline-concordant gestational weight gain.

Surveys of healthcare providers identified the following gaps requiring additional clinical support: increasing healthcare provider knowledge, improving skills and self-efficacy in discussing gestational weight gain and supporting women to make positive health behaviour changes in pregnancy.⁷⁴ System-related changes identified included flexibility in the time available for perinatal health appointments and changes in billing requirements. To address similar clinical gaps for healthcare providers in discussing weight gain in pregnancy, Alberta Health Services developed and evaluated an evidence-based, accredited online continuing medical education module which includes information about assessing, discussing and supporting healthy gestational weight gain in pregnant women. Interactive activities, self-testing and case scenarios help build capacity and provide opportunity to practice gained skills and knowledge. The module is available at <https://ecme.ucalgary.ca/programs/hpwg>.

Routine weighing: To date, RCTs of routine weighing of pregnant women during antenatal visits have not found routine weighing to be associated with significant impacts on gestational weight gain. First, a trial by Brownfoot et al.⁸⁹ of an intervention (regular weighing in antenatal clinics) compared with control (routine care) on gestational weight gain found no difference in the proportion of participants whose gestational weight gain was above, within or below the IOM recommended ranges among women with obesity. Second, McCarthy et al.⁹⁰ randomized 382 women with overweight or obesity and a singleton pregnancy, at less than 20 weeks' gestation and without diabetes to serial self-weighing and simple dietary advice (intervention $n=190$) or standard prenatal care (control $n=192$). There was no difference in gestational weight gain, for women with obesity, between the intervention and control groups (7.40 kg versus 8.77 kg).

Postpartum period

Background

The postpartum period is recognized to begin immediately following the birth of a baby, though its duration is not well defined. While many of the physiologic changes of pregnancy resolve by six weeks postpartum,^{91–93} several physiologic and psychological changes continue to exist for over a year postpartum, particularly if women continue to breastfeed.⁵² Thus, a period of one year after giving birth is generally considered as the working definition of the postpartum period.⁵²

A wide range of diverse and complex factors influence maternal weight after pregnancy. Higher gestational weight gain is recognized as a major risk factor for postpartum weight retention.^{12,47–49} The evidence is mixed with regard to the association between pre-pregnancy weight and postpartum weight retention, as several^{48,94–97} but not all^{98–100} studies have demonstrated that having a higher pre-pregnancy BMI is associated with higher

postpartum weight retention. Other factors, such as sleep patterns,¹⁰¹ psychological factors (e.g., depression, anxiety and stress symptoms),^{102–105} and maternal characteristics including age,^{48,106} ethnicity,^{48,99,107} parity,^{98,108} socioeconomic status,^{48,106,109} and educational attainment^{48,109} are associated with postpartum weight retention.

The findings that weight gain during pregnancy and the potential lack of, or limited, weight loss after pregnancy increases the risk of longer-term obesity or the risk moving into a higher BMI category are consistent across studies and worldwide. Wallace and colleagues³² for example, examined inter-pregnancy weight change using data from the Aberdeen Maternity and Neonatal Databank and identified that 86% of women who were initially categorized in the obesity BMI categories gained a further 9 kg by their third pregnancy. In Ireland, Mullaney⁵³ investigated the trajectory of postpartum weight change and found that 90% of women with obesity in early pregnancy retained obesity at four and nine months postpartum. In a prospective cohort study of Canadian women,⁹¹ BMI increased by approximately 1.5 kg/m² from pre-pregnancy to 10 to 12 weeks postpartum across all pre-pregnancy BMI categories. A population-based study of 58,534 Canadians who experienced successive pregnancies¹¹⁰ demonstrated that approximately 25% of women were ≥ 5 kg heavier at the beginning of a subsequent pregnancy, $\sim 9\%$ were heavier by 10 kg or more, and approximately 2% of women developed obesity after starting the previous pregnancy with a lower BMI. Retaining or gaining weight following pregnancy may also be associated with adverse pregnancy outcomes in future pregnancies. These adverse outcomes are described in the pre-conception and antenatal sections and in Table 1.

Many of the pregnancy-related contributors to gestational weight gain, such as fluid gained during pregnancy and increased organ size (e.g. of the uterus), are lost in the first six weeks postpartum. After this period, postpartum weight retention is mainly due to increased fat mass.⁹¹ Despite the adverse long-term health impact of postpartum weight retention, no detailed guidelines have been established to define cut-points for excess postpartum weight retention.¹¹¹ The dietary reference intakes established by the IOM (2005) 112 recommend a weight loss of approximately 0.8 kg/month in the first six months postpartum, but this amount of weight loss is not specifically for women with obesity. These guidelines are based on a review by Butte and Hopkinson (1998) of nine longitudinal studies conducted with affluent postpartum women, which showed that well-nourished women experience an average weight loss of 0.8 kg/month in the first six months postpartum. Weight stability is assumed after this period,¹¹³ and there is no consensus about the amounts or timelines for continued weight loss after the first six months postpartum.

For women who entered pregnancy with obesity, a postpartum weight loss of at least the amount gained during gestation should be achieved to not increase the risk of adverse pregnancy outcomes in future pregnancy.^{6,114} Sorbey and colleagues⁶ demonstrated that postpartum weight loss by more than two BMI units protects against the development of gestational diabetes in future

pregnancies among women with obesity. Postpartum weight gain increases the risk of gestational diabetes in the next pregnancy. McBain et al.¹¹⁴ also found that women in the overweight/obesity BMI categories were more likely to remain in the same BMI category in the next pregnancy.

Notably, evidence shows that women want an opportunity to discuss postpartum weight loss with their healthcare providers, and suggests that it should be a part of standard care.⁷³ This discussion could take place either during the six-week postnatal check-up or even during pregnancy in order for women to have the opportunity to start thinking about postpartum weight loss early on.⁷³ According to Ohlendorf (2012),¹¹⁵ 67% of postpartum women are attempting behavioural interventions or maintaining healthy behaviour at four weeks postpartum, and 84% at eight weeks postpartum. Most (82%) postpartum women with obesity reported that they planned to seek information about losing pregnancy weight through their physicians and midwives, as well as media.¹¹⁶

Clinical strategies and resources to achieve postpartum weight management for women with obesity

Considering the above evidence, the following clinical interventions should be considered to assist with weight management of postpartum women with obesity:

Physical activity and motivational interviewing: Physical activity and motivational interviewing during pregnancy have been shown to be beneficial to women with obesity in managing their postpartum weight loss. Claesson et al.²¹ undertook a study involving pregnant women with obesity who participated in a program to limit gestational weight gain by including regular physical activity (aqua aerobics class) and using motivational interviewing to identify and plan for individual behaviour change. Study participants had a significantly lower body weight at 10 to 12 weeks postpartum compared with controls (mean with standard deviation) (93.2 [13.32] kg versus 96.5 [14.48] kg; $p=0.037$). The intervention was delivered weekly during pregnancy and every six months during the two first years after childbirth. The effects of the intervention were assessed 12 and 24 months later.¹¹⁷ The mean weight change in the intervention group was -2.2 kg compared to +0.4 kg in the control group from early pregnancy to the follow-up 12 months after childbirth ($p=0.046$). More women in the intervention group showed sustained weight loss 24 months after delivery than women in the control group over the same time period ($p=0.034$). Women with obesity in the intervention group who gained less than 7 kg during pregnancy had a significantly lower weight than the controls at the 24-month follow-up ($p=0.018$).

Combined behavioural intervention: A comprehensive approach targeting both nutrition and exercise has been shown to be beneficial for women with obesity in managing postpartum weight loss. Vesco and colleagues¹² randomly assigned 114 pregnant women with obesity to either a group-based intervention or to a usual care control group to test the effectiveness of the intervention on maternal weight change from randomization to two

weeks postpartum. The intervention program included a combination of dietary and exercise recommendations, as well as the use of behavioural self-management techniques and attendance at weekly group meetings until delivery. Control participants received one-time dietary advice. The intervention group lost more weight at two weeks postpartum compared to control group (-2.6 vs +1.2 kg, mean difference of -3.8 kg; 95% CI [-5.9-1.7]; $p < 0.001$). However, the maternal weight-related benefits of this intervention did not persist at one year postpartum.¹¹⁸

Nascimento et al.¹⁶ conducted a systematic review and meta-analysis to test the effect of physical exercise strategies on weight loss in postpartum women with overweight or obesity (i.e., BMI ≥ 25 kg/m²). Though the results were not separated for women with BMI ≥ 25 kg/m², combined physical activity (with clear targets) and nutrition had greatest impacts on weight loss mean difference of -4.34 kg [95% CI -5.15 to -3.52, $I^2 = 0\%$] across all included studies.

Breastfeeding: Breastfeeding has been shown to improve a range of maternal and infant health outcomes, though the impacts on postpartum weight retention remain under study.³¹ However, studies have demonstrated that women with obesity have additional challenges with breastfeeding requiring individualized support.³¹ Specifically, a prospective cohort study²⁶ of 4231 postpartum women who were enrolled at the time of birth of their babies and were followed-up at three, 12, 24 and 48 months postpartum evaluated breastfeeding. Information was gathered on breastfeeding patterns and showed that the median duration of exclusive and total breastfeeding was decreased in the obesity group compared with the group with lower weight. Also, women with obesity have an increased probability of weaning their babies by three months postpartum compared with women with lower weight. Therefore, women with obesity are less likely to initiate and maintain breastfeeding^{26,119,120} compared with their counterparts with lower weight, and they may require additional supports on establishing and continuing breastfeeding.²⁶ Of note, the safety of pharmacotherapy agents for weight maintenance has not yet been established during breastfeeding; therefore these agents are generally avoided.²⁵

Gaps in the literature/future research for women with obesity over the reproductive years

1. Health outcomes in women with obesity (BMI ≥ 30 kg/m²): There remains a paucity of data specific to women with obesity (BMI ≥ 30 kg/m²), as many studies aggregate women with pre-pregnancy BMI ≥ 25 kg/m² into a single group for analysis for studies over the three reproductive periods. This is an important distinction, as women with a pre-pregnancy BMI ≥ 30 kg/m² may have different trajectories of gestational weight gain than those whose BMI is 25–29.9 kg/m².^{121,122}

2. Preconception weight management: There are emerging intervention studies on weight management preconception for women with obesity to improve fertility and obstetric outcomes. More data are needed, however, to examine how to implement

these interventions into clinical practice to engage a broader range of women with obesity and their healthcare providers.

3. Preconception counselling on pregnancy risks: Numerous observational studies have demonstrated that women with obesity have increased risks of several adverse pregnancy outcomes. Research is needed to identify effective counselling strategies to discuss pregnancy risks using a patient-centred approach.

4. Classes of obesity: The current IOM/Health Canada recommendations for gestational weight gain for women with obesity were not stratified by obesity classes, which may be associated with different obstetrical risks.⁵⁰ There remains a paucity of data regarding the impact of low weight gain or weight loss during pregnancy among women with different classes of obesity.¹²³

5. Low weight gain or weight loss during pregnancy: While healthcare providers commonly recommended gestational weight gain below guidelines for women with obesity, this advice is not consistent with the current recommendations. Research is needed to examine both the clinical outcomes and healthcare providers' reasons for providing this advice.¹²⁴

6. Patient-provider relationship: The impacts of the relationship between healthcare providers and their patients on gestational weight gain has not been extensively examined. Compared with women with lower weight, data demonstrate that providers asked fewer lifestyle questions and gave less lifestyle information to women with overweight and obesity. Point-of-care tools developed based on principles of patient-centredness and pregnant women's self-efficacy to manage gestational weight gain may be simple clinical tools, though these require further study.¹²²

7. Duration of pregnancy: The current recommendations for total gestational weight gain pertain to term pregnancies, typically defined as 37 to 41 completed weeks' gestation. However, evidence suggests that this group does not have homogeneous fetal outcomes.^{125,126} Future research needs to separate the independent effects of gestational weight gain from the independent effects of gestational age during this five-week period.

8. Postpartum weight management: In general, there remains a paucity of specific literature to guide women with obesity and their healthcare providers on the optimal timing, rate and amount of weight loss in the postpartum period and the impacts on clinically important health outcomes. In addition, more research is needed to examine factors that motivate and support women with obesity to engage in and maintain weight management strategies in the postpartum period.¹¹⁶

Correspondence:
guidelines@obesitynetwork.ca

References

1. Adane AA, Tooth LR, Mishra GD. Pre-pregnancy weight change and incidence of gestational diabetes mellitus : A finding from a prospective cohort study. *Diabetes Res Clin Pract.* 2017;124:72-80. doi:10.1016/j.diabres.2016.12.014
2. Savitri AI, Zuithoff P, Browne JL, et al. Does pre-pregnancy BMI determine blood pressure during pregnancy ? A prospective cohort study. *BMJ Open.* 2016;6(8):1-8. doi:10.1136/bmjopen-2016-011626
3. Ertel KA, Huang T, Mph SLR, et al. Perinatal weight and risk of prenatal and postpartum depressive symptoms. *Ann Epidemiol.* 2017;27(11):695-700.e1. doi:10.1016/j.annepidem.2017.10.007
4. Frederick IO, Rudra CB, Miller RS, et al. Adult weight change, weight cycling, and prepregnancy obesity in relation to risk of preeclampsia. *Epidemiology.* 2006;17(4):428-434. doi:10.1097/01.ede.0000221028.33245.0b
5. Rasmussen KM, Catalano PM, Yaktine AL. New guidelines for weight gain during pregnancy: what obstetrician/gynecologists should know. *Curr Opin Obstet Gynecol.* 2009;21(6):521-526. doi:10.1097/gco.0b013e328332d24e
6. Sorbye L, Klungsoyr K, Morken NH. Gestational diabetes mellitus and interpregnancy weight change : A population- based cohort study. *PLoS Med / Public Libr Sci.* 2017;14(8):1-19.
7. Krukowski RA, West D, Dicarolo M, et al. A Behavioral Intervention to Reduce Excessive Gestational Weight Gain. *Matern Child Health J.* 2017;21(3):485-491. doi:10.1007/s10995-016-2127-5
8. Agha M, Agha RA, Sandell J. Interventions to Reduce and Prevent Obesity in Pre- Conceptual and Pregnant Women : A Systematic Review and Meta-Analysis. *PLoS One.* 2014;9(5):e95132. doi:10.1371/journal.pone.0095132
9. Herring SJ, Cruice JF, Bennett GG, Rose MZ, Davey A, Foster GD. Preventing Excessive Gestational Weight Gain Among African American Women : A Randomized Clinical Trial. *Obesity.* 2016;24(1):30-36. doi:10.1002/oby.21240
10. Poston L, Bell R, Croker H, et al. Effect of a behavioural intervention in obese pregnant women (the UPBEAT study): a multicentre, randomised controlled trial. *Lancet Diabetes Endocrinol.* 2015;3(10):767-777. doi:10.1016/S2213-8587(15)00227-2
11. Haby K, Glantz A, Senior PD, et al. Mighty Mums – An antenatal healthcare intervention can reduce gestational weight gain in women with obesity. *Midwifery.* 2015;31(7):685-692. doi:10.1016/j.midw.2015.03.014
12. Vesco KK, Karanja N, King JC, et al. Efficacy of a Group-Based Dietary Intervention for Limiting Gestational Weight Gain among Obese Women : A Randomized Trial. *Obesity.* 2014;22(9):1989-1996. doi:10.1002/oby.20831
13. Vinter CA, J.S. Jørgensen, Ovesen P, Beck-Nielsen H, Skytthe A, Jensen DM. Metabolic effects of lifestyle intervention in obese pregnant women. Results from the randomized controlled trial "lifestyle in pregnancy" (LiP). *Diabet Med.* 2014;31(11):1323-1330. doi:10.1111/dme.12548
14. Petrella E, Malavolti M, Bertarini V, et al. Gestational weight gain in overweight and obese women enrolled in a healthy lifestyle and eating habits program. *J Matern Neonatal Med.* 2014;27(13):1348-1352. doi:10.3109/14767058.2013.858318
15. Harrison CL, Lombard CB, Strauss BJ, Teede HJ. Optimizing healthy gestational weight gain in women at high risk of gestational diabetes: A randomized controlled trial. *Obesity.* 2013;21(5):904-909. doi:10.1002/oby.20163
16. Nascimento SL, Pudwell J, Surita FG, Adamo KB, Smith GN. The effect of physical exercise strategies on weight loss in postpartum women: A systematic review and meta-analysis. *Int J Obes.* 2014;38(5):626-635. doi:10.1038/ijo.2013.183
17. Wolff S, Legarth J, Vangsgaard K, Toubro S, Astrup A. A randomized trial of the effects of dietary counseling on gestational weight gain and glucose metabolism in obese pregnant women. *Int J Obes.* 2008;32(3):495-501. doi:10.1038/sj.ijo.0803710
18. Nascimento SL, Surita FG, Parpinelli M, Siani S, Pinto e Silva JL. The effect of an antenatal physical exercise programme on maternal/perinatal outcomes and quality of life in overweight and obese pregnant women: A randomised clinical trial. *BJOG An Int J Obstet Gynaecol.* 2011;118(12):1455-1463. doi:10.1111/j.1471-0528.2011.03084.x
19. Barakat R, Pelaez M, Cordero Y, et al. Exercise during pregnancy protects against hypertension and macrosomia: Randomized clinical trial. *Am J Obstet Gynecol.* 2016;214(5):649.e1-649.e8. doi:10.1016/j.ajog.2015.11.039
20. Daly N, Farren M, McKeating A, O'Kelly R, Stapleton M, Turner MJ. A medically supervised pregnancy exercise intervention in obese women: A randomized controlled trial. *Obstet Gynecol.* 2017;130(5):1001-1010. doi:10.1097/AOG.0000000000002267
21. Claesson IM, Sydsjö G, Brynhildsen J, et al. Weight gain restriction for obese pregnant women: A case-control intervention study. *BJOG An Int J Obstet Gynaecol.* 2008;115(1):44-50. doi:10.1111/j.1471-0528.2007.01531.x
22. Chiswick C, Reynolds RM, Denison F, et al. Effect of metformin on maternal and fetal outcomes in obese pregnant women (EMPOWaR): A randomised, double-blind, placebo-controlled trial. *Lancet Diabetes Endocrinol.* 2015;3(10):778-786. doi:10.1016/S2213-8587(15)00219-3
23. Syngelaki A, Nicolaidis KH, Balani J, et al. Metformin versus placebo in obese pregnant women without diabetes mellitus. *N Engl J Med.* 2016;374(5):434-443. doi:10.1056/NEJMoa1509819
24. Dodd JM, Louise J, Deussen AR, et al. Effect of metformin in addition to dietary and lifestyle advice for pregnant women who are overweight or obese: the GRoW randomised, double-blind, placebo-controlled trial. *Lancet Diabetes Endocrinol.* 2019;7(1):15-24. doi:10.1016/S2213-8587(18)30310-3
25. Vitner D, Harris K, Maxwell C, Farine D. Obesity in pregnancy: a comparison of four national guidelines. *J Matern Neonatal Med.* 2019;32(15):2580-2590. doi:10.1080/14767058.2018.1440546
26. Castillo H, Santos IS, Matijasevich A. Maternal pre-pregnancy BMI, gestational weight gain and breastfeeding. *Eur J Clin Nutr.* 2016;70(4):431-436. doi:10.1038/ejcn.2015.232
27. Beckmann MM, Widmer T, Bolton E. Does preconception care work? *Aust New Zeal J Obstet Gynaecol.* 2014;54(6):510-514. doi:10.1111/ajo.12224

28. American College of Obstetricians and Gynecologists. Prepregnancy counseling. *Obstet Gynecol.* 2019;133:e78-e89. doi:10.1016/j.fertnstert.2018.12.003
29. Gesink Law DC, Macle hose RF, Longnecker MP. Obesity and time to pregnancy. *Hum Reprod.* 2007;22(2):414-420. doi:10.1093/humrep/del400
30. Paterson N, Sharma AM, Maxwell C, Greenblatt EM. Obesity-related health status is a better predictor of pregnancy with fertility treatment than body mass index: a prospective study. *Clin Obes.* 2016;6(4):243-248. doi:10.1111/cob.12149
31. Dutton H, Borengasser SJ, Gaudet LM, Barbour LA, Keely EJ. Obesity in pregnancy: optimizing outcomes for mom and baby. *Med Clin North Am.* 2018;102(1):87-106. doi:10.1016/j.physbeh.2017.03.040
32. Wallace JM, Bhattacharya S, Horgan GW. Weight change across the start of three consecutive pregnancies and the risk of maternal morbidity and SGA birth at the second and third pregnancy. *PLoS One.* 2017;12(6):e0179589. doi:10.1371/journal.pone.0179589
33. Liu Y, Dai W, Dai X, Li Z. Prepregnancy body mass index and gestational weight gain with the outcome of pregnancy: A 13-year study of 292,568 cases in China. *Arch Gynecol Obstet.* 2012;286(4):905-911. doi:10.1007/s00404-012-2403-6
34. Singh J, Huang C, Driggers RW, et al. The impact of pre-pregnancy body mass index on the risk of gestational diabetes The impact of pre-pregnancy body mass index on the risk of gestational diabetes. *J Matern Neonatal Med.* 2012;25(1):5-10. doi:10.3109/14767058.2012.626920
35. Fortner RT, Pekow P, Solomon CG, Markenson G, Chasan-taber L. Prepregnancy body mass index, gestational weight gain, and risk of hypertensive pregnancy among Latina women. *Am J Obstet Gynecol.* 2009;200(2):167-169. doi:10.1016/j.ajog.2008.08.021
36. Masho SW, Urban P, Cha S, Ramus R. Body Mass Index, Weight Gain, and Hypertensive Disorders in Pregnancy. *Am J Hypertens.* 2016;29(6):763-771. doi:10.1093/ajh/hpv184
37. Blondon M, Harrington LB, Boehlen F, Robert-ebadi H, Righini M, Smith NL. Pre-pregnancy BMI, delivery BMI, gestational weight gain and the risk of postpartum venous thrombosis. *Thromb Res.* 2016;145:151-156. doi:10.1016/j.thromres.2016.06.026
38. Macdonald SC, Bodnar LM, Himes KP, Hutcheon JA, Canada V, Sciences R. Patterns of gestational weight gain in early pregnancy and risk of gestational diabetes mellitus. *Epidemiology.* 2017;28(3):419-427. doi:10.1097/EDE.0000000000000629.Patterns
39. Hedderon MM, Gunderson EP, Ferrara A. Gestational Weight Gain and Risk of Gestational Diabetes Mellitus. *Obstet Gynecol.* 2010;115(3):597-604. doi:10.1097/AOG.0b013e3181cfce4f.Gestational
40. Cox Bauer CM, Bernhard KA, Greer DM, Merrill DC. Maternal and neonatal outcomes in obese women who lose weight during pregnancy. *J Perinatol.* 2016;36(4):278-283. doi:10.1038/jp.2015.202
41. Chandrasekaran S, Levine LD, Durnwald CP, Elovitz MA, Srinivas SK. Excessive weight gain and hypertensive disorders of pregnancy in the obese patient. *J Matern Neonatal Med.* 2015;28(8):964-968. doi:10.3109/14767058.2014.939624
42. Durst JK, Sutton ALM, Cliver SP, Tita AT, Biggio JR. Impact of gestational weight gain on perinatal outcomes in obese women. *Am J Perinatol.* 2016;33(09):849-855. doi:10.1055/s-0036-1579650
43. Faucher MA, Barger MK. Gestational weight gain in obese women by class of obesity and select maternal/newborn outcomes: A systematic review. *Women Birth J Aust Coll Midwives.* 2015;28(3):e70-e79. doi:10.1016/j.wombi.2015.03.006
44. Crane JMG, White J, Murphy P, Burrage L, Hutchens D. The Effect of Gestational Weight Gain by Body Mass Index on Maternal and Neonatal Outcomes. *J Obstet Gynaecol Canada.* 2009;31(1):28-35.
45. Lindsay KL, Brennan L, Rath A, Maguire OC, Smith T. Gestational weight gain in obese pregnancy : impact on maternal and foetal metabolic parameters and birthweight. *J Obstet Gynaecol (Lahore).* 2018;38(1):60-65. doi:10.1080/01443615.2017.1328670
46. Vesco KK, Dietz PM, Rizzo J, et al. Excessive gestational weight gain and postpartum weight retention among obese women. *Obstet Gynecol.* 2009;114(5):1069-1075. doi:10.1097/AOG.0b013e3181baeacf
47. Ashley-Martin J, Woolcott C. Gestational Weight Gain and Postpartum Weight Retention in a Cohort of Nova Scotian Women. *Matern Child Health J.* 2014;18(8):1927-1935. doi:10.1007/s10995-014-1438-7
48. Endres LK, Straub H, McKinney C, et al. Postpartum weight retention risk factors and relationship to obesity at 1 year. *Obstet Gynecol.* 2015;125(1):144-152. doi:10.1097/AOG.0000000000000565.Postpartum
49. Begum F, Colman I, McCargar LJ, Bell R. Gestational weight gain and early postpartum weight retention in a prospective cohort of Alberta women. *J Obstet Gynaecol Canada.* 2012;34(7):637-647. doi:10.1007/s10995-014-1438-7
50. Bogaerts A, Ameye L, Martens E, Devliege R. Weight loss in obese pregnant women and risk for adverse perinatal outcomes. *Obstet Gynecol.* 2015;125(3):566-575. doi:10.1097/AOG.0000000000000677
51. Oza-Frank R, Keim SA. Should Obese Women Gain Less Weight in Pregnancy Than Recommended? *Birth.* 2013;40(2):107-114. doi:10.1111/birt.12037
52. Walker LO, Wilging S. Rediscovering the "M" in "MCH": Maternal Health Promotion After Childbirth: THOUGHTS & OPINIONS. *J Obstet Gynecol Neonatal Nurs.* 2000;29(3):229-236. doi:10.1007/s00192-009-0937-3.Pelvic
53. Mullaney L, O'Higgins AC, Cawley S, Daly N, McCartney D, Turner MJ. Maternal weight trajectories between early pregnancy and four and nine months postpartum. *Public Health.* 2016;100(135):144-146. doi:10.1016/j.puhe.2016.02.017
54. Davies GAL, Maxwell C, McLeod L, et al. Obesity in pregnancy. *Int J Gynecol Obstet.* 2010;110(2):167-173. doi:10.1016/j.ijgo.2010.03.008
55. Molyneux E, Poston L, Khondoker M, Howard LM. Obesity, antenatal depression, diet and gestational weight gain in a population cohort study. *Arch Womens Ment Health.* 2016;19(5):899-907. doi:10.1007/s00737-016-0635-3
56. Lindsay KL, Heneghan C, McNulty B, Brennan L, McAuliffe FM. Lifestyle and Dietary Habits of an Obese Pregnant Cohort. *Matern Child Health J.* 2014;19(1):25-32. doi:10.1007/s10995-014-1491-2
57. McPhie S, Skouteris H, Hill B, Hayden M. Understanding gestational weight gain: The role of weight-related expectations and knowledge. *Aust New Zeal J Obstet Gynaecol.* 2015;55(1):21-26. doi:10.1111/ajo.12265
58. Holowko N, Mishra G, Koupil I. Social inequality in excessive gestational weight gain. *Int J Obes.* 2014;38(1):91-96. doi:10.1038/ijo.2013.62
59. Most J, Gilmore LA, Altazan AD, et al. Propensity for adverse pregnancy outcomes in African-American women may be explained by low energy expenditure in early pregnancy. *Am J Clin Nutr.* 2018;107(6):957-964. doi:10.1093/ajcn/nqy053
60. Guilloty NI, Soto R, Anzalota L, Rosario Z, Cordero JF, Palacios C. Diet, Pre-pregnancy BMI, and Gestational Weight Gain in Puerto Rican Women. *Matern Child Health J.* 2015;19(11):2453-2461. doi:10.1007/s10995-015-1764-4
61. Krukowski RA, Bursac Z, McGehee MA, West D. Exploring potential health disparities in excessive gestational weight gain. *J Women's Heal.* 2013;22(6):494-500. doi:10.1089/jwh.2012.3998
62. Walker LO, Hoke MM, Brown A. Risk factors for excessive or inadequate gestational weight gain among Hispanic women in a U.S.-Mexico border state. *JOGNN - J Obstet Gynecol Neonatal Nurs.* 2009;38(4):418-429. doi:10.1111/j.1552-6909.2009.01036.x
63. Bogaerts A, Van den Bergh B, Nuyts E, Martens E, Witters I, Devlieger R. Socio-demographic and obstetrical correlates of pre-pregnancy body mass index and gestational weight gain. *Clin Obes.* 2012;2(5-6):150-159. doi:10.1111/cob.12004
64. Weeks A, Halili L, Liu RH, Deonandan R, Adamo KB. Gestational weight gain counselling gaps as perceived by pregnant women and new mothers: Findings from the electronic maternal health survey. *Women and Birth.* 2019. doi:10.1016/j.wombi.2019.02.005

65. Washington Cole KO, Gudzone KA, Bleich SN, et al. Influence of the 5A's Counseling Strategy on Weight Gain during Pregnancy: An Observational Study. *J Women's Heal.* 2017;26(10):1123-1130. doi:10.1089/jwh.2016.6115
66. Canadian Obesity Network. 5As of Healthy Pregnancy Weight Gain.; 2014. https://obesitycanada.ca/wp-content/uploads/2018/02/CON_HealthyPregnancy_book_15_final.pdf.
67. Stotland NE, Haas JS, Brawarsky P, Jackson RA, Fuentes-Afflick E, Escobar GJ. Body mass index, provider advice, and target gestational weight gain. *Obstet Gynecol.* 2005;105(3):633-638. doi:10.1097/01.AOG.0000152349.84025.35
68. Mercado A, Marquez B, Abrams B, Phipps MG, Wing RR, Phelan S. Where Do Women Get Advice About Weight, Eating, and Physical Activity During Pregnancy? *J Womens Health (Larchmt).* 2017;26(9):951-956. doi:10.1089/jwh.2016.6078
69. Yeo SA, Walker JS, Caughey MC, Ferraro AM, Asafu-Adjei JK. What characteristics of nutrition and physical activity interventions are key to effectively reducing weight gain in obese or overweight pregnant women? A systematic review and meta-analysis. *Obes Rev.* 2017;18(4):385-399. doi:10.1111/obr.12511
70. Deputy NP, Sharma AJ, Kim SY, Olson CK. Achieving Appropriate Gestational Weight Gain: The Role of Healthcare Provider Advice. *J Women's Heal.* 2018;27(5):552-560. doi:10.1089/jwh.2017.6514
71. Criss S, Oken E, Guthrie L, Hivert MF. A qualitative study of gestational weight gain goal setting. *BMC Pregnancy Childbirth.* 2016;16(1):317. doi:10.1186/s12884-016-1118-2
72. Allen-Walker V, Mullaney L, Turner MJ, et al. How do women feel about being weighed during pregnancy? A qualitative exploration of the opinions and experiences of postnatal women. *Midwifery.* 2017;49:95-101. doi:10.1016/j.midw.2016.12.006
73. Nikolopoulos H, Mayan M, Maclsaac J, Miller T, Bell RC. Women's perceptions of discussions about gestational weight gain with healthcare providers during pregnancy and postpartum: A qualitative study. *BMC Pregnancy Childbirth.* 2017;17(1):97. doi:10.1186/s12884-017-1257-0
74. Morris J, Nikolopoulos H, Berry T, et al. Healthcare providers' gestational weight gain counselling practises and the influence of knowledge and attitudes: A cross-sectional mixed methods study. *BMJ Open.* 2017;7(11):e018527. doi:10.1136/bmjopen-2017-018527
75. Ferraro ZM, Boehm KS, Gaudet LM, Adamo KB. Counseling about gestational weight gain and healthy lifestyle during pregnancy: Canadian maternity care providers' self-evaluation. *Int J Womens Health.* 2013;5:629-636. doi:10.2147/IJWH.S49422
76. Phelan S, Phipps MG, Abrams B, Darroch F, Schaffner A, Wing RR. Practitioner advice and gestational weight gain. *J Women's Heal.* 2011;20(4):585-591. doi:10.1089/jwh.2010.2316
77. Stengel MR, Kraschnewski JL, Hwang SW, Kjerulff KH, Chuang CH. "What My Doctor Didn't Tell Me": Examining Healthcare Provider Advice to Overweight and Obese Pregnant Women on Gestational Weight Gain and Physical Activity. *Women's Heal Issues.* 2012;22(6):e535-e540. doi:10.1016/j.whi.2012.09.004
78. Emery RL, Benno MT, Salk RH, Kolko RP, Levine MD. Healthcare provider advice on gestational weight gain: uncovering a need for more effective weight counselling. *J Obstet Gynaecol (Lahore).* 2018;38(7):916-921. doi:10.1080/01443615.2018.1433647
79. Fieldwick D, Smith A, Paterson H. General practitioners and gestational weight management. *J Obstet Gynaecol (Lahore).* 2019;39(4):485-491. doi:10.1080/01443615.2018.1530739
80. Piccinini-Vallis H. Clinician self-efficacy in initiating discussions about gestational weight gain. *Can Fam Physician.* 2017;63(7):e341-e349.
81. Stewart M, Brown JB, Weston W, McWhinney IR, McWilliam CL, Freeman T. Patient-centered medicine: transforming the clinical method. CRC Press. 2013. doi:10.5334/ijc.116
82. Renault KM, Nørgaard K, Nilas L, et al. The Treatment of Obese Pregnant Women (TOP) study: A randomized controlled trial of the effect of physical activity intervention assessed by pedometer with or without dietary intervention in obese pregnant women. *Am J Obstet Gynecol.* 2014;210(2):134.e1-134.e9. doi:10.1016/j.ajog.2013.09.029
83. Dodd J., Grivell RM, Deussen A, Eames AJ. Metformin for women who are obese during pregnancy versus standard care for improving maternal and infant outcomes. *Cochrane Database Syst Rev.* 2018;(7). doi:10.1002/14651858.CD010564.pub2. www.cochranelibrary.com
84. Saha S, Beach MC. The impact of patient-centered communication on patients' decision making and evaluations of physicians: A randomized study using video vignettes. *Patient Educ Couns.* 2011;84(3):386-392. doi:10.1016/j.pec.2011.04.023
85. Michie S, Miles J, Weinman J. Patient-centredness in chronic illness: What is it and does it matter? *Patient Educ Couns.* 2003;51(3):197-206. doi:10.1016/S0738-3991(02)00194-5
86. Leske S, Strodl E, Hou XY. Patient-practitioner relationships desired by overweight/obese adults. *Patient Educ Couns.* 2012;89(2):309-315. doi:10.1016/j.pec.2012.07.002
87. Little P, Everitt H, Williamson I, et al. Observational study of effect of patient centredness and positive approach on outcomes of general practice consultations. *Br Med J.* 2001;323(7318):908-911. doi:10.1136/bmj.323.7318.908
88. Lindberg SM, DeBoth A, Anderson CK. Effect of a Best Practice Alert on Gestational Weight Gain, Health Services, and Pregnancy Outcomes. *Matern Child Health J.* 2016;20(10):2169-2178. doi:10.1007/s10995-016-2052-7
89. Brownfoot FC, Davey MA, Kornman L. Routine weighing to reduce excessive antenatal weight gain: A randomised controlled trial. *BJOG An Int J Obstet Gynaecol.* 2016;123(2):254-261. doi:10.1111/1471-0528.13735
90. McCarthy EA, Walker SP, Ugoni A, Lappas M, Leong O, Shub A. Self-weighing and simple dietary advice for overweight and obese pregnant women to reduce obstetric complications without impact on quality of life: A randomised controlled trial. *BJOG An Int J Obstet Gynaecol.* 2016;123(6):965-973. doi:10.1111/1471-0528.13919
91. Schmitt NM, Nicholson WK, Schmitt J. The association of pregnancy and the development of obesity - Results of a systematic review and meta-analysis on the natural history of postpartum weight retention. *Int J Obes.* 2007;31(11):1642-1651. doi:10.1038/sj.ijo.0803655
92. Gunderson EP, Abrams B, Selvin S. Does the pattern of postpartum weight change differ according to pregravid body size? *Int J Obes.* 2001;25(6):853-862. doi:10.1038/sj.ijo.0801631
93. Østbye T, Peterson BL, Krause KM, Swamy GK, Lovelady CA. Predictors of postpartum weight change among overweight and obese women: Results from the active mothers postpartum study. *J Women's Heal.* 2012;21(2):215-222. doi:10.1089/jwh.2011.2947
94. Linné Y, Dye L, Barkeling B, Rössner S. Weight development over time in parous women - The SPAWN study: 15 Years follow-up. *Int J Obes.* 2003;27(12):1516-1522. doi:10.1038/sj.ijo.0802441
95. Kirkegaard H, Stovring H, Rasmussen KM, Abrams B, Sorensen TIA, Nohr EA. How do pregnancy-related weight changes and breastfeeding relate to maternal weight and BMI-adjusted waist circumference 7 y after delivery? Results from a path analysis. *Am J Clin Nutr.* 2013;99(2):312-319. doi:10.3945/ajcn.113.067405.INTRODUCTION
96. Ketterl TG, Dundas NJ, Roncaioli SA, Littman AJ, Phipps AI. Association of Pre-pregnancy BMI and Postpartum Weight Retention Before Second Pregnancy, Washington State, 2003-2013. *Matern Child Health J.* 2018;22(9):1339-1344. doi:10.1007/s10995-018-2514-1
97. Nohr EA, Vaeth M, Baker JL, Sørensen TIA, Olsen J, Rasmussen KM. Combined associations of prepregnancy body mass index and gestational weight gain with the outcome of pregnancy. *Am J Clin Nutr.* 2008;87(6):1750-1759. doi:10.3945/ajcn.2008.26939
98. Hollis JL, Crozier SR, Inskip HM, et al. Modifiable risk factors of maternal postpartum weight retention: An analysis of their combined impact and potential opportunities for prevention. *Int J Obes.* 2017;41(7):1091-1098. doi:10.1038/ijo.2017.78
99. Waage CW, Falk RS, Sommer C, et al. Ethnic differences in postpartum weight retention: A Norwegian cohort study. *BJOG An Int J Obstet Gynaecol.* 2016;123(5):699-708. doi:10.1111/1471-0528.13321

100. Rong K, Yu K, Han X, et al. Pre-pregnancy BMI, gestational weight gain and postpartum weight retention: A meta-analysis of observational studies. *Public Health Nutr.* 2015;18(12):2172-2182. doi:10.1017/S1368980014002523
101. Gunderson EP, Rifas-Shiman SL, Oken E, et al. Association of fewer hours of sleep at 6 months postpartum with substantial weight retention at 1 year postpartum. *Am J Epidemiol.* 2007;167(2):178-187. doi:10.1093/aje/kwm298
102. Pedersen P, Baker JL, Henriksen TB, et al. Influence of psychosocial factors on postpartum weight retention. *Obesity.* 2011;19(3):639-646. doi:10.1038/oby.2010.175
103. Whitaker K, Young-Hyman D, Vernon M, Wilcox S. Maternal stress predicts postpartum weight retention. *Matern Child Health J.* 2014;18(9):2209-2217. doi:10.1007/s10995-014-1470-7
104. Phillips J, King R, Skouteris H. A conceptual model of psychological predictors of postpartum weight retention. *J Reprod Infant Psychol.* 2012;30(3):278-288. doi:10.1080/02646838.2012.717265
105. Herring SJ, Rich-Edwards JW, Oken E, Rifas-Shiman SL, Kleinman KP, Gillman MW. Association of postpartum depression with weight retention 1 year after childbirth. *Obesity.* 2008;16(6):1296-1301. doi:10.1038/oby.2008.71
106. Olson CM, Strawderman MS, Hinton PS, Pearson TA. Gestational weight gain and postpartum behaviors associated with weight change from early pregnancy to 1 y postpartum. *Int J Obes.* 2003;27(1):117-127. doi:10.1038/sj.ijo.0802156
107. Parker JD, Abrams B. Differences in postpartum weight retention between black and white mothers. *Obstet Gynecol.* 1993;81(5 Pt 1):768-774.
108. Hill B, Bergmeier H, McPhie S, et al. Is parity a risk factor for excessive weight gain during pregnancy and postpartum weight retention? A systematic review and meta-analysis. *Obes Rev.* 2017;18(7):755-764. doi:10.1111/obr.12538
109. Shrewsbury VA, Robb KA, Power C, Wardle J. Socioeconomic differences in weight retention, weight-related attitudes and practices in postpartum women. *Matern Child Health J.* 2009;13(2):231-240. doi:10.1007/s10995-008-0342-4
110. Hutcheon JA, Chapinal N, Bodnar LM, Lee L. The INTERGROWTH-21st gestational weight gain standard and interpregnancy weight increase: A population-based study of successive pregnancies. *Obesity.* 2017;25(6):1122-1127. doi:10.1002/oby.21858
111. Adegboye ARA, Linne YM. Diet or exercise, or both, for weight reduction in women after childbirth. *Cochrane Database Syst Rev.* 2013;(7). doi:10.1002/14651858.CD005627.pub3
112. Institute of Medicine. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids. Washington, DC: The National Academies Press. <https://doi.org/10.17226/10490>. doi:10.1016/S0002-8223(02)90346-9
113. Trumbo P, Schlicker S, Yates AA, Poos M. Food and Nutrition Board of the Institute of Medicine, The National Academies. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids. *J Am Diet Assoc.* 2002;102(11):1621-1630. doi:10.1016/S0002-8223(02)90346-9
114. McBain RD, Dekker GA, Clifton VL, Mol BW, Grzeskowiak LE. Impact of inter-pregnancy BMI change on perinatal outcomes: a retrospective cohort study. *Eur J Obstet Gynecol Reprod Biol.* 2016;205:98-104. doi:10.1016/j.ejogrb.2016.07.487
115. Ohlendorf JM. Stages of Change in the Trajectory of Postpartum Weight Self-Management. *J Obstet Gynecol Neonatal Nurs.* 2012;41(1):57-70. doi:10.1111/j.1552-6909.2011.01323.x
116. Ohlendorf JM, Weiss ME, Ryan P. Weight-management information needs of postpartum women. *MCN Am J Matern Nurs.* 2012;37(1):56-63. doi:10.1097/NMC.0b013e31823851ee
117. Claesson IM, Sydsjö G, Brynhildsen J, et al. Weight after childbirth: A 2-year follow-up of obese women in a weight-gain restriction program. *Acta Obstet Gynecol Scand.* 2011;90(1):103-110. doi:10.1111/j.1600-0412.2010.01016.x
118. Vesco KK, Leo MC, Karanja N, et al. One-year postpartum outcomes following a weight management intervention in pregnant women with obesity. *Obesity.* 2016;24(10):2042-2049. doi:10.1002/oby.21597
119. Britz SE, McDermott KC, Pierce CB, Blomquist JL, Handa VL. Changes in maternal weight 5-10 years after a first delivery. *Women's Heal.* 2012;8(5):513-519. doi:10.2217/whe.12.35
120. Turckson R, Bel S, Galjaard S, Devlieger R. Maternal obesity and breastfeeding intention, initiation, intensity and duration: A systematic review. *Matern Child Nutr.* 2014;10(2):166-183. doi:10.1111/j.1740-8709.2012.00439.x
121. Jarman M, Yuan Y, Pakseresht M, Shi Q, Robson PJ, Bell RC. Patterns and trajectories of gestational weight gain: a prospective cohort study. *C Open.* 2016;4(2):E338-E345. doi:10.9778/cmajo.20150132
122. Piccinini-Vallis H, Vallis M. Curbing excess gestational weight gain in primary care: Using a point-of-care tool based on behavior change theory. *Int J Womens Health.* 2018;10:609-615. doi:10.2147/IJWH.S172346
123. Kapadia MZ, Park CK, Beyene J, Giglia L, Maxwell C, McDonald SD. Weight loss instead of weight gain within the guidelines in obese women during pregnancy: A systematic review and meta-analyses of maternal and infant outcomes. *PLoS One.* 2015;10(7):e0132650. doi:10.1371/journal.pone.0132650
124. Delgado A, Stark LM, Macri CJ, Power ML, Schulkin J. Provider and Patient Knowledge and Views of Office Practices on Weight Gain and Exercise during Pregnancy. *Am J Perinatol.* 2018;35(2):201-208. doi:10.1055/s-0037-1606582
125. Fleischman AR, Oinuma M, Clark SL. Rethinking the definition of "term pregnancy." *Obstet Gynecol.* 2010;116(1):136-139. doi:10.1097/AOG.0b013e3181e24f28
126. Spong CY. Defining "term" pregnancy: Recommendations from the defining "term" pregnancy workgroup. *JAMA - J Am Med Assoc.* 2013;309(23):2445-2446. doi:10.1001/jama.2013.6235