

Bariatric Surgery: Selection & Pre-Operative Workup

Stephen Glazer MDⁱ, Laurent Biertho MDⁱⁱ

- i) Department of Internal Medicine, University of Toronto
- ii) Department of Surgery, Laval University

Cite this Chapter

Glazer S, Biertho L. Canadian Adult Obesity Clinical Practice Guidelines: Bariatric Surgery: Selection & Pre-Operative Workup. Available from: <https://obesitycanada.ca/guidelines/preop>. Accessed [date].

Update History

Version 1, August 4, 2020. The Canadian Adult Obesity Clinical Practice Guidelines are a living document, with only the latest chapters posted at obesitycanada.ca/guidelines.

KEY MESSAGES FOR HEALTHCARE PROVIDERS

- Criteria for selection of appropriate candidates for bariatric surgery have been established to minimize surgical complications and to maximize the benefit of these important and limited procedures.
- The pre-operative workup should evaluate a patient's medical, nutritional, mental and functional health status.
- Special attention should be given to the care of patients living with type 2 diabetes (T2DM) who are considering bariatric surgery to minimize complications from uncontrolled diabetes in the peri-operative period.
- Because of the risks of post-operative complications associated with tobacco use, cessation prior to bariatric surgery is mandatory and should be maintained lifelong.
- In patients living with severe obesity, bariatric surgery, in combination with behavioural interventions, is a more effective option for long-term weight loss and control of chronic conditions, such as T2DM, hypertension, sleep apnea and dyslipidemia, as well as other conditions associated with increased adiposity.

RECOMMENDATIONS

- We suggest a comprehensive medical and nutritional evaluation be completed and nutrient deficiencies corrected in candidates for bariatric surgery (Level 4, Grade D).^{1,2}
- We suggest screening for and treatment of obstructive sleep apnea in people seeking bariatric surgery (Level 4, Grade D).^{4,5}
- Pre-operative smoking cessation can minimize post-operative complications (Level 2a, Grade B).³

KEY MESSAGES FOR PEOPLE LIVING WITH OBESITY

- Bariatric surgery is the beginning of a life-long journey. You should educate yourself about the necessary changes required to optimize your long-term outcomes for a healthier life.
- Before surgery you will be asked to perform several investigations, such as blood work and cardiac or pulmonary testing, to ensure that you are ready and safe for surgery.
- If you are at high risk for obstructive sleep apnea, you may be asked to undergo a sleep study to determine if you have significant sleep apnea.
- A current or recent history of smoking or nicotine consumption puts you at risk of complications after bariatric surgery. Smoking cessation is required before surgery, and must be maintained for life.
- You may be given a low-calorie diet two to three weeks before surgery in order to shrink your liver size and make your surgery easier.
- If you are living with diabetes, you will have to follow your blood sugars very closely and obtain instructions on how to adjust your diabetes medications while on the low-calorie diet prior to bariatric surgery.
- Because changes in the absorption of some medications may occur with certain bariatric surgical procedures, you may be asked to change either the type or preparation of the medication you are currently taking.

Selection of appropriate patients for bariatric surgery

Extensive preparation of patients prior to bariatric surgery is required. Potential candidates for bariatric surgery undergo multi-disciplinary evaluation and optimization of their medical, mental, nutritional and functional health to assess their eligibility and safety to proceed with surgery. Further medical evaluations may include cardiac, respiratory, metabolic, gastrointestinal and sleep apnea testing. Once adequate evaluation and preparation have been undertaken, the patient may proceed for bariatric surgery if stable. It is recommended that patients engage in behavioural interventions prior to bariatric surgery and maintain those behavioural changes after bariatric surgery as well.

Bariatric surgery is indicated in patients greater than 18 years of age with a body mass index (BMI) of 35 kg/m², who have at least one major obesity-related complication including type 2 diabetes (T2DM), hypertension, hyperlipidemia, pseudotumor cerebri, obesity hypoventilation syndrome or Pickwickian syndrome, debilitating arthritis, non-alcoholic fatty liver disease or non-alcoholic steatohepatitis, coronary artery disease, severe reflux or obstructive sleep apnea.¹ Bariatric surgery is also indicated for patients with a BMI of 40 kg/m² independent of the presence of obesity-related complications.⁶ Bariatric surgery may also provide therapeutic benefit in patients with a BMI between 30 and 34.9 kg/m² who have been refractory to non-surgical attempts at weight loss with obesity-related complications, especially T2DM. The indications for surgery in these patients may be predicated heavily on the obesity-related complications that are present and weighing the durable response to bariatric surgery compared with existing medical interventions.^{7,8}

Adolescent candidates who have a BMI greater than or equal to 35 kg/m² may be considered for a bariatric procedure if they possess

additional significant complications, including T2DM, significant obstructive sleep apnea with an apnea-hypopnea index measured at 15 or higher, pseudotumor cerebri or severe non-alcoholic steatohepatitis (NASH). In addition, consideration may be given to those adolescent candidates with a BMI of greater than or equal to 40 kg/m² with other complications, such as hypertension, insulin resistance or glucose intolerance, significant impairment with their quality of life or activities of daily living, hyperlipidemia or obstructive sleep apnea.⁹

General considerations for bariatric surgery candidates

Potential surgical candidates should have a history of identifiable attempts at the medical management of obesity. All patients must be committed to engage in the educational process involved in preparing for bariatric surgery as well as adhering to the necessary long-term follow-up, both from a nutritional and medical perspective. Appropriate surveillance and treatment of potential long-term nutritional deficiencies as well as assessment and intervention for their obesity-related complications and mental health is crucial for long-term success.

Patients with unstable psychiatric illness, malignancy or other diseases associated with decreased life expectancy, substance abuse or an inability to adhere to long-term follow-up may be considered inappropriate candidates for surgery due to a high risk of short- and long-term complications.

Advanced patient age is not an absolute contraindication to bariatric surgery. The outcomes and complication rates for patients greater than 60 years of age appear to be comparable to those of a younger population, regardless of the surgical procedure performed.¹⁰

Non-alcoholic steatohepatitis NASH is common in patients living with obesity and can lead to liver cirrhosis. The risk of bariatric surgery in patients carefully selected with Child-Pugh Class A liver cirrhosis is not prohibitive, but caution and additional surveillance should be undertaken as their overall risk for perioperative complications and mortality is increased.^{11,12} In addition, non-alcoholic steatohepatitis is the third most common indication for transplantation of the liver and, with growing incidence, NASH is becoming an increasingly frequent cause for liver transplantation.¹³⁻¹⁵

The morbidity and mortality rate of bariatric surgery before or after liver transplantation is increased but remains acceptable in tertiary care centres.¹⁶

Predictors of successful post-operative weight loss

Behavioural changes

Exercise in conjunction with obesity management programs has been demonstrated to improve weight loss and body composition. Thirty minutes per day (150 minutes per week) of moderate-intensity exercise after bariatric surgery is associated with a 3.6 kg additional weight loss compared with individuals who do not exercise after bariatric surgery.¹⁷

In preparation for bariatric surgery, candidates may benefit from implementing health behaviour change interventions. Interventions include extensive education on nutrition and the need for exercise, and behavioural strategies for successful weight loss and weight maintenance, exploring topics such as self-monitoring, mindless eating and goal setting. However, bariatric surgical patients subjected to a behavioural intervention program for six months prior to bariatric surgery experienced no significant difference in the degree of their weight loss at 24 months after surgery compared with control subjects.¹⁸

Patients' concurrent eating patterns may play a significant role in the effect of post-operative weight loss. In addition, past traumas may also impact upon weight loss outcomes post-bariatric surgery. The presence of a history of binge eating disorder or a previous history of sexual abuse did not seem to predict difference in weight loss outcomes at two years post-bariatric surgery, but this area remains controversial and requires further study.¹⁹

Pre-operative weight loss

Weight reduction is associated with improvement of cardiac risk factors and the associated complications of obesity. As little as 5-10% weight loss can improve cardiovascular risk factors and reduce complications, such as hypertension, hyperlipidemia, T2DM, visceral fat and hepatic steatosis, as well as liver volume. High-protein diets produce rapid weight loss, provide adequate satiety, reduce lean body mass loss and decrease reduction in the resting energy expenditure.²⁰ There does not appear to be any compromise in immune function or any effect on wound healing with the use of high-protein diets.

Failure to achieve adequate post-operative weight loss outcomes is often attributed to pre-operative psychosocial characteristics and eating behaviours, as well as poor adherence to the recommended post-operative nutrition plan after bariatric surgery.²¹ Past weight management attempts pre-operatively may reflect a patient's ability to follow a strict medical nutrition plan and exercise program post-operatively.²¹

Pre-operative weight stability is desired for patients prior to bariatric surgery. Pre-operative weight loss may decrease the difficulty of performing bariatric surgery, minimize blood loss, improve short-term weight loss and short-term complications, as well as decrease operative time.²²⁻²⁴

Patients who achieved $\geq 5\%$ excess body weight loss pre-operatively experienced more substantial weight loss at one year compared with those with less pre-operative weight loss.²⁵ However, longer-term studies reviewing pre-operative weight loss outcomes did not confer any advantage at four years with respect to weight loss outcomes.²⁶

Overall, there is no compelling evidence mandating weight loss prior to bariatric surgery for the long-term efficacy of weight management, but rather only for the technical simplicity of the surgery.²¹

Pre-surgical medical nutrition therapy

Many pre-operative protocols include the use of a liquid-based, low-calorie diet for two to three weeks prior to surgery.²⁷ Low-calorie diets in Canada provide around 900 calories per day, consisting of low carbohydrates, high protein and low-fat intake.

Bariatric surgical patients are requested to consume very low-calorie meal supplements in the form of commercially available protein shakes totalling 650-900 kcal/day for two to three weeks prior to surgery. A low-calorie diet which is low in carbohydrates may cause reduced liver volume by up to 19% and a 17% reduction in visceral adipose tissue.^{27,28}

A systematic review found that very low-calorie diets (VLCDs) are effective at reducing body weight and hepatic volume prior to surgery. These diets do not reduce intra-operative or post-surgical risks.²⁷ They do result in a surgeon's perception that the difficulty of performing the operation is decreased.²²

The effect of pre-operative weight management also provides additional motivation to the patient in preparation for bariatric surgery. However, adherence and compliance with pre-operative meal supplements may be poor and occasionally not well tolerated. In addition, meal supplements may be expensive. Pre-operative weight loss with a two-week VLCD may approach approximately 6 kg.²⁹

Pre-operative patients taking a VLCD using Optifast® 800 kcal per day for two weeks prior to surgery were compared with patients ingesting the same caloric intake from a normal diet. There was a non-significant difference in the operating time, but there was

a higher surgeon-perceived scale of difficulty in control patients compared with patients who received a VLCD prior to surgery. The 30-day complication rate was higher in control patients, including wound GI and deep hemorrhage, infection, dehiscence and anastomotic leak.²² In another study, pre-operative weight loss of $\geq 8\%$ excess weight in patients following a VLCD for four weeks prior to surgery was associated with decreased hospital length of stay and greater three-month and one-year post-operative weight loss compared with those who did not follow a VLCD.^{23,30} Neither the major complication rate nor conversion rates were affected by the degree of pre-operative weight loss.²³

Risk assessment prior to bariatric surgery

Nutritional evaluation

Limited high-quality evidence has reviewed pre-operative malnutrition status in patients seeking bariatric surgery. Nonetheless, observational studies have indicated that patients living with obesity have a higher risk for inadequate nutritional status^{2,31-33} and malnutrition.^{31,34,35} Pre-operative evaluation and collaborative support from a registered dietitian is recommended for all patients considering bariatric surgery.^{1,2}

A large, multi-center retrospective observational study (n = 106,577) found that ~6% of patients undergoing bariatric surgery were malnourished and had increased risk of death or serious morbidity and 30-day readmission rates.³⁴ This study also found that > 10% weight loss prior to surgery was associated with nine-times higher rates of DSM (Diagnostic and Statistical Manual of Mental Disorders) in patients with mild malnutrition, and 10-times higher DSM in those with severe malnutrition.³⁴ Similarly, a retrospective cohort study concluded that 32% of the patient cohort (n = 533) had malnutrition prior to surgery.³⁵ Higher BMI was associated with increased risk for malnutrition, and post-operative nausea and vomiting were associated with pre-operative malnutrition.³⁵

Prevalence of pre-operative micronutrient deficiencies is also higher in patients with obesity.³⁶ Pre-operative lab work should include complete blood count, creatinine, iron panel, vitamin D, calcium, albumin and vitamin B12. Fasting plasma glucose, hemoglobin A1C, lipid panel and liver enzymes can also be measure at the same time, as part of the evaluation of obesity-related metabolic complications. Vitamin A, parathormone, phosphate, zinc, selenium and copper levels can be assessed more selectively, due to cost considerations. Pre-operative optimization of micronutrient levels prior to surgery, specifically levels of vitamin D, vitamin B12 and iron, is recommended.² Pre-operative multi-vitamin complex with vitamin B1 is usually started at least one month prior to surgery. Of note, patients taking proton pump inhibitors and/or metformin have a higher prevalence of vitamin B12 deficiency.²

Smoking and nicotine use

Smoking and nicotine cessation should be recommended for all patients undergoing bariatric surgery. Surgical centres recommend

that smoking cessation should be achieved for at least six weeks up to six months prior to bariatric surgery to ensure adequate extinction of consumption.³⁷

Nicotine contributes to ulcer development by potentiating acid and pepsin secretion, increases bile salt reflux, increases H. pylori infection risk and diminishes prostaglandin synthesis, mucosal blood flow and gastric mucus.³⁸

Cigarette smoking may be associated with increased risk of marginal ulceration and may increase the risk of stricture formation. Complications related to smoking exposure on average may occur post-operatively around 14 months.³⁹

Smoking post-bariatric surgery may also be associated with pneumonia and post-operative complications with subsequent surgeries involving body contour surgery and mastopexy.^{3,40-44}

Pre-operative investigations

The pre-operative evaluation of a bariatric surgical candidate is similar to those patients considered for similar-risk surgery, with some caveats. Bariatric surgical candidates may have additional factors which may make their clinical evaluation more challenging. Many patients referred for bariatric surgery may have a low or exceptionally low functional capacity.

Cardiac evaluation: Exercise tolerance is a predictor for surgical outcomes; symptom-limited stairclimbing, for example, is predictive of post-operative cardiopulmonary complications in patients undergoing high-risk surgery.⁴⁵ In addition, obesity is an independent risk factor for cardiovascular disease. Many patients with obesity may also convey symptoms of shortness of breath or chest pain, and the etiology of these symptoms may be varied. Furthermore, the physical examination in patients living with obesity may be challenging in that physical findings such as distant heart sounds or the evaluation of the jugular venous pressure may be difficult to obtain.

Patients undergoing bariatric surgery may have abnormalities in their electrocardiogram (ECG). An abnormal ECG in patients undergoing bariatric surgery may be independently associated with a higher likelihood of a complicated post-operative course, including the need for post-operative intensive care (ICU) admissions.⁴⁶ Obesity may also be associated with changes in cardiac morphology, including left ventricular hypertrophy, diastolic dysfunction or left ventricular dysfunction. Bariatric surgery is associated with a decrease in left ventricular mass index, left ventricular end-diastolic volume, improvement in diastolic dysfunction and systolic function and left atrial diameter.⁴⁷

Cardiac echo is not typically performed during pre-operative evaluations, but may be required to detect suspected left or right ventricular dysfunction, valvular heart disease or pulmonary hypertension. The visualization of the echocardiographic images is often suboptimal, and echocardiographic contrast agents improve

the ability to identify endocardial borders and to assess ventricular wall motion abnormalities. A radionuclide angiography (RNA) multigated acquisition (MUGA) scan may also be helpful in the assessment of ejection fraction. Cardiac nuclear stress testing may be required to investigate chest pain. In patients with BMI > 30 kg/m², an abnormal nuclear stress test was associated with a higher annual cardiac event rate, cardiac death or death from any cause.⁴⁸ Computed tomographic coronary angiography may be difficult with increased body habitus due to depth-dependent spatial resolution. In addition, computerized tomography scanner tables may be limited by weight restriction. The gold standard for cardiac evaluation remains coronary angiography, which was not associated with an increase in minor or major complication rates in patients living with obesity.⁴⁹

Pulmonary evaluation: The respiratory system is also significantly affected in patients with an elevated BMI. Impairment in pulmonary function due to restriction in lung volumes and abnormalities in respiratory mechanics resulting in increased work of breathing may detrimentally affect the respiratory status of bariatric surgical candidates.^{50,51}

Sleep apnea: It is not uncommon for people living with obesity to experience sleep-related disorders, which may result in significant respiratory, cardiovascular and neuropsychiatric conditions.⁵² Obstructive sleep apnea, one type of a sleep-related disorder, is either the complete cessation of airflow (apnea) or the significant reduction of airflow (hypopnea) measured during sleep. The presence of obstructive sleep apnea has been associated with premature death,⁵³ motor vehicle accidents,⁵⁴ hypertension,⁵⁵ coronary artery disease and cerebrovascular accidents,⁵⁶ nocturnal cardiac arrhythmias⁵⁷ and T2DM.⁵⁸ In addition, obstructive sleep apnea has been associated with a significant increase in the incidence of sudden death from all cardiac causes.⁵⁹ Obstructive sleep apnea has been demonstrated to affect the white matter within the limbic system, pons, frontal, temporal and parietal cortices, and projections connecting the cerebellum, which are key areas for brain function and memory.⁶⁰

The incidence of obstructive sleep apnea in middle-aged adults in the Wisconsin Sleep Cohort Study was approximately 24% in males and 9% in females.⁶¹ In patients undergoing bariatric surgery, the prevalence of obstructive sleep apnea can be ≥ 90%, with clinically significant sleep apnea underdiagnosed in as many as 50% of these patients.^{4,5} The prevalence of obstructive sleep apnea is much higher in males compared with female bariatric surgical candidates.⁶² The recognition of the diagnosis of obstructive sleep apnea is even more challenging as there may be no correlation between the severity of obstructive sleep apnea and BMI.⁶³

Bariatric surgery results in improvement or resolution of a patient's obstructive sleep apnea.⁶⁴ Obstructive sleep apnea in the bariatric population is believed to be caused by excess fat deposition in or around the neck, causing the patient's upper airway passages to collapse.⁶⁵ Patients undergoing bariatric surgery who have obstructive sleep apnea may have a higher complication rate, which may include a prolonged hospital stay, the occurrence of

thromboembolic phenomena, the need for reintervention and an increased 30-day mortality rate.⁶⁶ In addition, the presence of obstructive sleep apnea may be associated with more challenging intubations and increased ICU admissions.^{67,68}

The gold standard in the diagnosis of obstructive sleep apnea is a level 1 polysomnogram (PSG).⁶⁹ Resources for overnight in-laboratory PSG testing can be limited and expensive.⁷⁰ As a result, alternative methods for identifying patients at risk for obstructive sleep apnea have been established. Utilizing screening questionnaires, including the [STOP-Bang Questionnaire](#), [Berlin Questionnaire](#) and [Epworth Sleepiness Scale](#), has become routine for patients undergoing surgery. To identify bariatric surgical patients with clinically significant obstructive sleep apnea, relying on subjective screening questionnaires may fail to identify at risk patients.^{5,71} While some advocate that all bariatric surgical patients should be subjected to formal PSG, such resources are limited and costly.⁶³

The STOP-Bang or the Berlin Questionnaire may not be effective tools for detecting moderate- or high-risk patients for obstructive sleep apnea in patients undergoing bariatric surgery, and so more effective tools should be considered or developed.⁵ There may also be no significant correlation between the Epworth Sleepiness Score and severity of obstructive sleep apnea.⁷² There should be a high clinical suspicion for the presence of obstructive sleep apnea in bariatric patients undergoing surgery, with a low threshold for subjecting the patient to a formal PSG. In addition, the use of continuous positive airway pressure immediately post-operatively after gastric bypass is safe and should be administered if deemed clinically indicated.⁷³

Bariatric surgery is beneficial in improving obstructive sleep apnea.⁷⁴ However, despite significant weight loss post-operatively, bariatric surgical patients may have persistence of moderate to severe obstructive sleep apnea at one year post-operatively.⁷⁵

The long-term relationship between weight loss and sleep apnea is complex. There should be hypervigilance for recurrence of the presence of obstructive sleep apnea in patients previously diagnosed with obstructive sleep apnea, as reappearance can occur in the absence of weight regain.^{76,77}

Endoscopy: There exists controversy in the utility of pre-operative endoscopy in patients undergoing bariatric surgery. Most patients who undergo endoscopy have normal or non-clinically significant findings. The consideration for performing endoscopy should be individualized based on symptoms, risk factors and type of procedure being considered for each patient.⁷⁸ Patients considering sleeve gastrectomy who have dyspepsia, reflux, dysphagia or symptoms suggestive of foregut pathology, as well as those on chronic anti-acid therapy, should undergo pre-operative endoscopy to rule out the presence of hiatal hernia, esophagitis or Barrett's esophagus or other diseases such as peptic ulcer disease and tumours.⁷⁹⁻⁸³ Screening for *H. pylori* can be performed at the time of endoscopy. The incidence of *H. pylori* in patients with obesity planning to undergo bariatric surgery is variable, ranging between 15% and 85%.^{83,84} *H. pylori* may be implicated in the

development of gastritis, peptic ulcer and gastric carcinoma.^{85,86} Screening for *H. pylori* is recommended for this reason in patients undergoing gastric bypass procedure. The benefits for patients undergoing other types of surgeries is controversial.⁸⁷ Screening investigations for malignancy should also be considered prior to bariatric surgery due to the association between certain malignancies and obesity. This may include screening colonoscopy for malignancy in patients 50 years of age and over, and mammography and Pap smears in appropriate candidates.⁸⁸⁻⁹⁰

Risk of thromboembolism: The 90-day incidence of venous thromboembolism after bariatric surgery is 0.42%. Although uncommon, the clinical consequences can be devastating. Up to 40% of peri-operative deaths may be attributed to pulmonary embolism. This remains one of the most common causes of peri-operative death along with myocardial infarction and sepsis from anastomotic leak.⁹¹ Prophylaxis with low molecular weight heparin to prevent thromboembolism post-operatively after gastric bypass is common practice. The prophylactic use of an inferior vena cava filter is no longer recommended, even in patients at high risk of pulmonary embolism, as it is associated with increased risk of post-operative deep vein thrombosis and overall mortality without decreasing the risk of pulmonary embolism.⁹²

Other consideration: Patients with obesity are at increased risk for several gastrointestinal, hepatobiliary and intra-abdominal processes. Evaluation of the bariatric surgical patient with an abdominal ultrasound is not routinely recommended, except in those patients requiring an investigation for symptomatic biliary disease and elevated liver enzymes or non-alcoholic fatty liver disease.⁹³

Following bariatric surgery, decreases in bone density may be observed due to bone loss. Mixed restrictive and malabsorptive procedures, such as Roux-en-Y gastric bypass and biliopancreatic diversion, increase risk for bone fractures. Sleeve gastrectomy may result in bone loss to a lesser degree. Bone loss post-bariatric surgery may be attributed to many factors, including nutritional factors, skeletal unfolding, calciotropic hormone abnormalities, body and bone marrow fat changes and changes in gut hormones. Baseline bone density evaluation may be considered prior to bariatric surgery and two years post-bariatric surgery depending on risk factors, which include post-menopausal women, older men, patients with prior fragility fractures or a family history of osteoporosis. Vitamin D and parathyroid hormone levels may be obtained pre-operatively in the screening for patients at risk for metabolic bone disease.⁹⁴

Medication considerations

Prior to bariatric surgery, patients need to receive instructions and general precautions surrounding their medications. Avoidance of aspirin may be required prior to surgery in patients taking it for primary prevention. In addition, anti-inflammatory agents must be discontinued prior to surgery. The utilization of these agents post-operatively will depend on their indication, risk tolerance and the surgical procedure. Chronic use of non-steroidal anti-inflammatory

drugs is contraindicated for Roux-en-Y Gastric bypass, due to the risk of anastomotic ulcer. Anti-platelet and anti-coagulant medication will also require cessation prior to surgery. In some cases, bridging anti-coagulation may be necessary. The use of direct oral anti-coagulants whose absorption is not dependent on low pH conditions may be considered for anti-coagulation post-sleeve gastrectomy. The efficacy of direct oral anti-coagulants post Roux-en-Y gastric bypass is uncertain, and hence vitamin K antagonists like warfarin remain the preferred oral agent for anti-coagulation.⁹⁵ Patients should be made aware of the need to switch to vitamin K antagonists after bypass procedures.

Immune-modulating medication used in the treatment of connective tissue and inflammatory disorders, skin disorders and immune-mediated gastrointestinal diseases may need to be held prior to surgery as well as post-operatively for a period of time, at the discretion of the prescribing specialist.

Long-acting release medications may need to be converted after bariatric surgery to short-acting preparations. Medications dependent upon absorption or an acid environment within the stomach and upper gastrointestinal tract may need to be re-evaluated, as well.⁹⁶⁻⁹⁸ Certain medications may need to be crushed, while encapsulated formulations may need to be opened in the early post-operative period. A comprehensive pharmacologic consultation prior to surgery should be considered.

Women taking estrogen therapy in the form of oral contraception should discontinue their medication four weeks prior to surgery while post-menopausal women may discontinue hormone replacement therapy three weeks prior to surgery.⁶⁶

Pre-operative management of patients living with diabetes

In preparation for bariatric surgery, blood glucose readings of patients living with T2DM should be optimized. Re-evaluation of the comprehensive care plan should be undertaken, as well as re-evaluation of their dietary intake, activity level and the status of other pharmacotherapy. Existing guidelines recommend targets for diabetic glycemic control with the hope of improving bariatric surgical outcomes. Suggested targets include A1C less than 7%, fasting blood sugar level less than 6.1 mmol/L and a two-hour postprandial blood sugar of 7.7 mmol/L. However, there is limited data guiding the management of glycemic control in bariatric surgery, both pre-operatively and post-operatively. With other surgeries, such as orthopedic and colorectal surgery, an elevated A1C pre-operatively may be associated with prolonged length-of-stay post-operatively and worsened post-operative outcomes.⁹⁹⁻¹⁰³

Elevated blood sugars prior to surgery may be associated with increased post-operative complications, decreased weight loss and less resolution of the patient's T2DM.¹⁰⁴

Randomized controlled trials suggest that neither intensive management of patients' glycemic control three months prior to Roux-en-Y

gastric bypass, nor intensive control of glycemia in the first two weeks post-operatively, resulted in better hemoglobin A1C levels one year after surgery.¹⁰⁵

Less aggressive blood glucose targets may be required prior to bariatric surgery, as often bariatric surgical patients with T2DM have high insulin resistance, resulting in suboptimal control.¹⁰⁶ While the patient with T2DM is on a controlled VLCD in preparation for bariatric surgery, re-evaluation of their diabetic medications should be undertaken, as a patient's requirements for medication to control blood sugars may be significantly altered. For diet-controlled patients with T2DM, no significant interventions are required.

Avoiding sulfonylureas and meglitinides while on VLCD should be considered to avoid hypoglycemia. In addition, sodium glucose co-transporter 2 inhibitors should be stopped while on a VLCD due to the risk of diabetic ketoacidosis.¹⁰⁷ Alpha-glucosidase, alpha-amylase enzyme inhibitors and thiazolidinediones may be stopped during this period of time. In addition, dipeptidyl peptidase-4 inhibitors and glucagon-like peptide 1 (GLP-1) agonists may also be held, especially with adequate blood sugar control.

Insulin requirements while on meal replacement therapy in preparation for surgery drop dramatically. Intermediate and long-acting insulin require a decreased dose, often by 50%, and short-acting insulins require significant re-adjustment, as well.^{108,109} Frequent blood sugar

monitoring is required while on VLCDs in preparation for bariatric surgery. Symptomatic hypoglycemia is treated in the usual fashion.¹¹⁰

While on meal replacement therapy, there may be an increased risk of intravascular volume depletion. Close observation of a patient's volume status, electrolytes and kidney function is a prudent and cautious approach.¹¹¹ Diuretics should be dose adjusted or held. In addition, close evaluation of blood pressure readings is required, and adjustment to anti-hypertensive medication may be necessary.¹¹² Patients with hypertension on blood pressure medication and concurrent meal replacement therapy should be educated about the possibility of developing orthostatic hypotension.

Individuals on meal replacement therapy and warfarin may also require closer observation of their international normalized ratio level.^{113,114}

Conclusion

Bariatric surgery is a life-altering and effective obesity management intervention. Several considerations must be undertaken to prepare a potential candidate for surgery. The patient's medical, mental, nutritional and functional health should be evaluated prior to surgery. Once adequate evaluation, preparation and optimization have been undertaken, establishing an acceptable peri-operative risk profile, the patient may proceed with bariatric surgery.

Downloaded from: <https://obesitycanada.ca/guidelines/preop>

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/) (CC BY-NC-ND 4.0)

The summary of the Canadian Adult Obesity Clinical Practice Guideline is published in the [Canadian Medical Association Journal](https://www.cma.ca/), and contains information on the full methodology, management of authors' competing interests, a brief overview of all recommendations and other details. More detailed guideline chapters are published on the Obesity Canada website at www.obesitycanada.ca/guidelines.

Correspondence:
guidelines@obesitynetwork.ca

References

1. Mechanick JI, Youdim A, Jones DB, et al. Clinical practice guidelines for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient-2013 update: Cosponsored by American association of clinical endocrinologists, the obesity society, and American society of. *Obesity*. 2013;21(SUP-PL. 1):S1-S27. doi:10.1002/oby.20461
2. Parrott J, N RD, Frank L, et al. American Society for Metabolic and Bariatric Surgery Integrated Health Nutritional Guidelines for the Surgical Weight Loss Patient 2016 Update : Micronutrients. *Surg Obes Relat Dis*. 2017;13(5):727-741. doi:10.1016/j.soard.2016.12.018
3. Myers K, Hajek P, Hinds C, McRobbie H. Stopping smoking shortly before surgery and postoperative complications: A systematic review and meta-analysis. *Arch Intern Med*. 2011;171(11):983-989. doi:10.1001/archinternmed.2011.97
4. Gasa M, Salord N, Fortuna AM, et al. Obstructive sleep apnoea and metabolic impairment in severe obesity. *Eur Respir J*. 2011;38(5):1089-1097. doi:10.1183/09031936.00198810
5. Glazer SA, Erickson AL, Crosby RD, Kieda J, Zawisza A, Deitel M. The Evaluation of Screening Questionnaires for Obstructive Sleep Apnea to Identify High-Risk Obese Patients Undergoing Bariatric Surgery. *Obes Surg*. 2018;28(11):3544-3552. doi:10.1007/s11695-018-3547-7
6. National Institutes of Health. National Heart, Lung and Bl. clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. 1998;158(17):1855-1867. doi:10.1001/archinte.158.17.1855
7. Busetto L, Dixon J, De Luca M, Shikora S, Pories W, Angrisani L. Bariatric surgery in class I obesity. *Obes Surg*. 2014;24(4):487-519. doi:10.1007/s11695-014-1214-1
8. ASMBS Clinical Issues Committee. Bariatric surgery in class I obesity (body mass index 30-35 kg/m²). *Surg Obes Relat Dis*. 2013;9(1):e1-e10. doi:10.1016/j.soard.2012.09.002
9. Pratt JSA, Browne A, Browne NT, Bruzoni M, Cohen M, Desai A, Inge T, Linden BC, Mattar SG, Michalsky M, Podkameni D, Reichard KW, Stanford FC, Zeller MH, Zitsman J. ASMBS pediatric metabolic and bariatric surgery guidelines, 2018. *Surg Obes Relat Dis*. 2018;14(7):882-901.

10. Giordano S, Vvictorzon M. Bariatric surgery in elderly patients: A systematic review. *Clin Interv Aging*. 2015;10:1627-1635. doi:10.2147/CIA.S70313
11. Shimizu H, Phuon V, Maia M, et al. Bariatric surgery in patients with liver cirrhosis. *Surg Obes Relat Dis*. 2013;9(1):1-6. doi:10.1016/j.soard.2012.07.021
12. Jan A, Narwaria M, Mahawar KK. A Systematic Review of Bariatric Surgery in Patients with Liver Cirrhosis. *Obes Surg*. 2015;25(8):1518-1526. doi:10.1007/s11695-015-1727-2
13. Agopian VG, Kaldas FM, Hong JC, et al. Liver transplantation for nonalcoholic steatohepatitis: The new epidemic. *Ann Surg*. 2012;256(4):624-633. doi:10.1097/SLA.0b013e31826b4b7e
14. Mandell MS, Zimmerman M, Campsen J, Kam I. Bariatric surgery in liver transplant patients: Weighing the evidence. *Obes Surg*. 2008;18(12):1515-1516. doi:10.1007/s11695-008-9712-7
15. Angulo P. Nonalcoholic fatty liver disease and liver transplantation. *Liver Transplant*. 2007;12(4):523-534. doi:10.1002/lt
16. Lazzati A, Iannelli A, Schneck AS, et al. Bariatric Surgery and Liver Transplantation: a Systematic Review a New Frontier for Bariatric Surgery. *Obes Surg*. 2015;25(1):134-142. doi:10.1007/s11695-014-1430-8
17. Egberts K, Brown WA, Brennan L, O'Brien PE. Does exercise improve weight loss after bariatric surgery? A systematic review. *Obes Surg*. 2012;22(2):335-341. doi:10.1007/s11695-011-0544-5
18. Kalarichian MA, Marcus MD, Courcoulas AP, Cheng Y, Levine MD. Preoperative lifestyle intervention in bariatric surgery: a randomized clinical trial. *Surg Obes Relat Dis*. 2016;12(1):180-187. doi:10.1016/j.physbeh.2017.03.040
19. Fujioka K, Yan E, Wang HJ, Li Z. Evaluating preoperative weight loss, binge eating disorder, and sexual abuse history on Roux-en-Y gastric bypass outcome. *Surg Obes Relat Dis*. 2008;4(2):137-143. doi:10.1016/j.soard.2008.01.005
20. Goldstein DJ. Beneficial health effects of modest weight loss. *Int J Obes Relat Metab Disord J Int Assoc Study Obes*. 1992;16(6):397-415.
21. Ríos B. *Integrated Health in Bariatric Surgery*.; 2015.
22. Van Nieuwenhove Y, Dambrauskas Z, Campillo-Soto A, et al. Preoperative very low-calorie diet and operative outcome after laparoscopic gastric bypass: A randomized multicenter study. *Arch Surg*. 2011;146(11):1300-1305. doi:10.1001/archsurg.2011.273
23. Alami RS, Morton JM, Schuster R, et al. Is there a benefit to preoperative weight loss in gastric bypass patients? A prospective randomized trial. *Surg Obes Relat Dis*. 2007;3(2):141-145. doi:10.1016/j.soard.2006.11.006
24. Cassie S, Menezes C, Birch DW, Shi X, Karmali S. Effect of preoperative weight loss in bariatric surgical patients: a systematic review. *Surg Obes Relat Dis*. 2011;7(6):767-768. doi:10.1016/j.soard.2011.09.003
25. Solomon H, Liu GY, Alami R, Morton J, Curet MJ. Benefits to Patients Choosing Preoperative Weight Loss in Gastric Bypass Surgery: New Results of a Randomized Trial. *J Am Coll Surg*. 2009;208(2):241-245. doi:10.1016/j.jamcollsurg.2008.09.028
26. Becouarn G, Topart P, Ritz P. Weight loss prior to bariatric surgery is not a pre-requisite of excess weight loss outcomes in obese patients. *Obes Surg*. 2010;20(5):574-577. doi:10.1007/s11695-010-0083-5
27. Holderbaum M, Casagrande DS, Sussenbach S, Buss C. Effects of very low calorie diets on liver size and weight loss in the preoperative period of bariatric surgery: a systematic review. *Surg Obes Relat Dis*. 2018;14(2):237-244. doi:10.1016/j.soard.2017.09.531
28. Benjaminov O, Beglaibter N, Gindy L, et al. The effect of a low-carbohydrate diet on the nonalcoholic fatty liver in morbidly obese patients before bariatric surgery. *Surg Endosc Other Interv Tech*. 2007;21(8):1423-1427. doi:10.1007/s00464-006-9182-8
29. Schouten R, van der Kaaden I, van 't Hof G, Feskens PGBM. Comparison of Preoperative Diets Before Bariatric Surgery: a Randomized, Single-Blinded, Non-inferiority Trial. *Obes Surg*. 2016;26(8):1743-1749. doi:10.1007/s11695-015-1989-8
30. Hutcheon DA, Hale AL, Ewing JA, et al. Short-Term Preoperative Weight Loss and Postoperative Outcomes in Bariatric Surgery. *J Am Coll Surg*. 2018;226(4):514-524. doi:10.1016/j.jamcollsurg.2017.12.032
31. Peterson LA, Cheskin LJ, Furtado M, et al. Malnutrition in Bariatric Surgery Candidates: Multiple Micronutrient Deficiencies Prior to Surgery. *Obes Surg*. 2016;26(4):833-838. doi:10.1007/s11695-015-1844-y
32. Sánchez A, Rojas P, Basfi-fer K, et al. Micronutrient Deficiencies in Morbidly Obese Women Prior to Bariatric Surgery. *Obes Surg*. 2016;26(2):361-368. doi:10.1007/s11695-015-1773-9
33. Dagan SS, Zelber-Sagi S, Webb M, et al. Nutritional Status Prior to Laparoscopic Sleeve Gastrectomy Surgery. *Obes Surg*. 2016;26(9):2119-2126. doi:10.1007/s11695-016-2064-9
34. Fieber JH, Sharoky CE, Wirtalla C, Williams NN, Dempsey DT, Kelz RR. The malnourished patient with obesity: a unique paradox in bariatric surgery. *J Surg Res*. 2018;232:456-463. doi:10.1016/j.jss.2018.06.056
35. Major P, Malczak P, Wysocki M, et al. Bariatric patients' nutritional status as a risk factor for postoperative complications, prolonged length of hospital stay and hospital readmission: A retrospective cohort study. *Int J Surg*. 2018;56:210-214. doi:10.1016/j.ijssu.2018.06.022
36. Gehr S, Kern B, Peters T, Christofiel-Courtin C, Peterli R. Fewer nutrient Deficiencies after laparoscopic sleeve gastrectomy (LSG) than after Laparoscopic Roux-Y-gastric bypass (LRYGB)-a prospective study. *Obes Surg*. 2010;20(4):447-453. doi:10.1007/s11695-009-0068-4
37. Devlin CA, Smeltzer SC. Temporary perioperative tobacco cessation: A literature review. *AORN J*. 2017;106(5):415-423. doi:10.1016/j.aorn.2017.09.001
38. Endoh K, Leung F. Effects of Smoking and Nicotine on the Gastric Mucosa: A Review of Clinical and Experimental Evidence. *Gastroenterology* 1994;107:884-878
39. Fakuluj A, Fong B, Abdemur A, Szomstein S, Rosenthal RJ. PL-114: Prevalence and implications of cigarette smoking in morbidly obese patients undergoing bariatric surgery. *Surg Obes Relat Dis*. 2010;6(3):S6. doi:10.1016/j.soard.2010.03.019
40. Felix EL, Kettelle J, Mobley E, Swartz D. Perforated marginal ulcers after laparoscopic gastric bypass. *Surg Endosc*. 2008;22(10):2128-2132. doi:10.1007/s00464-008-9996-7
41. Gupta PK, Gupta H, Kaushik M, et al. Predictors of pulmonary complications after bariatric surgery. *Surg Obes Relat Dis*. 2012;8(5):574-581. doi:10.1016/j.soard.2011.04.227
42. Gravante G, Araco A, Sorge R, Araco F, Delogu D, Cervelli V. Wound infections in body contouring mastopexy with breast reduction after laparoscopic adjustable gastric bandings: The role of smoking. *Obes Surg*. 2008;18(6):721-727. doi:10.1007/s11695-007-9252-6
43. Coupaye M, Sabaté JM, Castel B, et al. Predictive factors of weight loss 1 year after laparoscopic gastric bypass in obese patients. *Obes Surg*. 2010;20(12):1671-1677. doi:10.1007/s11695-010-0159-2
44. Garb J, Welch G, Zagarins S, Kuhn J, Romanelli J. Bariatric surgery for the treatment of morbid obesity: a meta-analysis of weight loss outcomes for laparoscopic adjustable gastric banding and laparoscopic gastric bypass. *Obes Surg*. 2009;19(10):1447-1455. doi:10.1007/s11695-009-9927-2
45. Girish M, Trayner E, Dammann O, Pinto-Plata V, Celli B. Symptom-limited stair climbing as a predictor of postoperative cardiopulmonary complications after high-risk surgery. *Chest*. 2001;120(4):1147-1151. doi:10.1378/chest.120.4.1147
46. Gonzalez R, Bowers SP, Venkatesh KR, Lin E, Smith CD. Preoperative factors predictive of complicated postoperative management after Roux-en-Y gastric bypass for morbid obesity. *Surg Endosc Other Interv Tech*. 2003;17(12):1900-1904. doi:10.1007/s00464-003-8810-9
47. Aggarwal R, Harling L, Efthimiou E, Darzi A, Athanasiou T, Ashrafian H. The Effects of Bariatric Surgery on Cardiac Structure and Function: a Systematic Review of Cardiac Imaging Outcomes. *Obes Surg*. 2016;26(5):1030-1040. doi:10.1109/ciced.2018.8592188

48. Elhendy A, Schinkel AFL, Van Domburg RT, et al. Prognostic stratification of obese patients by stress 99mTc- tetrofosmin myocardial perfusion imaging. *J Nucl Med.* 2006;47(8):1302-1306.
49. McNulty PH, Ettinger SM, Field JM, et al. Cardiac catheterization in morbidly obese patients. *Catheter Cardiovasc Interv.* 2002;56(2):174-177. doi:10.1002/ccd.10186
50. Aguiar IC, Freitas WR, Santos IR, et al. Obstructive sleep apnea and pulmonary function in patients with severe obesity before and after bariatric surgery: A randomized clinical trial. *Multidiscip Respir Med.* 2014;9(1):43. doi:10.1186/2049-6958-9-43
51. Wei YF, Wu HD. Candidates for bariatric surgery: Morbidly obese patients with pulmonary dysfunction. *J Obes.* 2012. doi:10.1155/2012/878371
52. Shahi B, Praglowski B, Deitel M. Sleep-related Disorders in the Obese. *Obes Surg.* 1992;2(2):157-168. doi:10.1381/096089292765560358
53. Hallowell PT, Stellato TA, Schuster M, et al. Potentially life-threatening sleep apnea is unrecognized without aggressive evaluation. *Am J Surg.* 2007;193(3):364-367. doi:10.1016/j.amjsurg.2006.09.022
54. Marshall NS, Wong KK, Liu PY, Cullen SR, Knuiaman MW, Grunstein RR. Sleep apnea as an independent risk factor for all-cause mortality: the Busselton Health Study. *Sleep.* 2008;31(8):1079-1085.
55. Terán-Santos J, Jiménez-Gómez A, Cordero-Guevara J. The association between sleep apnea and the risk of traffic accidents. *N Engl J Med.* 1999;340(11):847-851. doi:10.1056/NEJM199903183401104
56. Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Invest.* 1995;96(4):1897-1904. doi:10.1172/JCI118235
57. Parra O, Arboix A, Bechich S, et al. Time course of sleep-related breathing disorders in first-ever stroke or transient ischemic attack. *Am J Respir Crit Care Med.* 2000;161(2 I):375-380. doi:10.1164/ajrccm.161.2.9903139
58. Mehra R, Benjamin EJ, Shahar E, et al. Association of nocturnal arrhythmias with sleep-disordered breathing: The sleep heart health study. *Am J Respir Crit Care Med.* 2006;173(8):910-916. doi:10.1164/rccm.200509-14420C
59. Tasali E, Mokhlesi B, Van Cauter E. Obstructive sleep apnea and type 2 diabetes: Interacting epidemics. *Chest.* 2008;133(2):496-506. doi:10.1378/chest.07-0828
60. Macey PM, Kumar R, Woo MA, Valladares EM, Yan-Go FL, Harper RM. Brain structural changes in obstructive sleep apnea. *Sleep.* 2008;31(7):967-977. doi:10.1017/CBO9781139088268
61. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The Occurrence of Sleep-Disordered Breathing among Middle-Aged Adults. *N Engl J Med.* 1993;328(17):1230-1235. doi:10.1056/NEJM199304293281704
62. Peromaa-Haavisto P, Tuomilehto H, Kössi J, et al. Prevalence of Obstructive Sleep Apnoea Among Patients Admitted for Bariatric Surgery. A Prospective Multicentre Trial. *Obes Surg.* 2016;26(7):1384-1390. doi:10.1007/s11695-015-1953-7
63. O'Keefe T, Patterson EJ. Evidence Supporting Routine Polysomnography before Bariatric Surgery. *Obes Surg.* 2004;14(1):23-26. doi:10.1381/096089204772787248
64. Quintas-Neves M, Preto J, Drummond M. Assessment of bariatric surgery efficacy on obstructive sleep apnea (OSA). *Rev Port Pneumol (English Ed.)* 2016;22(6):331-336. doi:10.1016/j.rppnen.2016.05.006
65. Fritscher LG, Canani S, Mottin CC, et al. Bariatric Surgery in the Treatment of Obstructive Sleep Apnea in Morbidly Obese Patients. *Respiration.* 2007;74(6):647-652. doi:10.1159/000107736
66. Longitudinal Assessment of Bariatric Surgery (LABS) Consortium. Perioperative safety in the longitudinal assessment of bariatric surgery. *N Engl J Med.* 2009;361(5):445-454. doi:10.1056/NEJMoa0901836
67. Gupta RM, Parvizi J, Hanssen AD, Gay PC. Postoperative complications in patients with obstructive sleep apnea syndrome undergoing hip or knee replacement: a case-control study. *Mayo Clin Proc.* 2001;76(9):897-905.
68. Chesson J, Ferber RA, Fry JM, et al. Practice parameters for the indications for polysomnography and related procedures. *Sleep.* 1997;20(6):406-422. doi:10.1093/sleep/20.6.406
69. Gentil B, De Larminat JM, Boucherez C, Lienhart A, Chou HC, Wu TL. Difficult intubation and obstructive sleep apnoea syndrome. *Br J Anaesth.* 1994;72(3):368.
70. Nagappa M, Liao P, Wong J, et al. Validation of the stop-bang questionnaire as a screening tool for obstructive sleep apnea among different populations: A systematic review and meta-Analysis. *PLoS One.* 2015;10(12):e0143697. doi:10.1371/journal.pone.0143697
71. Sharkey KM, Orff HJ, Tosi C, Harrington D, Roye GD, Millman RP. Subjective sleepiness and daytime functioning in bariatric patients with obstructive sleep apnea. *Sleep Breath.* 2013;17(1):267-274. doi:10.1007/s11325-012-0685-3
72. Serafini FM, Anderson WM, Rosemurgy AS, Strait T, Murr MM. Clinical predictors of sleep apnea in patients undergoing bariatric surgery. *Obes Surg.* 2001;11(1):28-31. doi:10.1381/096089201321454079
73. Ramirez A, Lalor PF, Szomstein S, Rosenthal RJ. Continuous positive airway pressure in immediate postoperative period after laparoscopic Roux-en-Y gastric bypass: is it safe? *Surg Obes Relat Dis.* 2009;5(5):544-546. doi:10.1016/j.soard.2009.05.007
74. Sarkhosh K, Switzer NJ, El-Hadi M, Birch DW, Shi X, Karmali S. The impact of bariatric surgery on obstructive sleep apnea: a systematic review. *Obes Surg.* 2013;23(3):414-423. doi:10.1016/B978-0-12-420168-2.00040-5
75. Peromaa-Haavisto P, Tuomilehto H, Kössi J, et al. Obstructive sleep apnea: the effect of bariatric surgery after 12 months. A prospective multicenter trial. *Sleep Med.* 2017;35(July):85-90. doi:10.1016/j.sleep.2016.12.017
76. Sampol G, Muñoz X, Sagalés MT, et al. Long-term efficacy of dietary weight loss in sleep apnoea/hypopnoea syndrome. *Eur Respir J.* 1998;12(5):1156-1159. doi:10.1183/09031936.98.12051156
77. Pillar G, Peled R, Lavie P. Recurrence of sleep apnea without concomitant weight increase 7.5 years after weight reduction surgery. *Chest.* 1994;106(6):1702-1704. doi:10.1378/chest.106.6.1702
78. Parikh M, Liu J, Vieira D, et al. Preoperative Endoscopy Prior to Bariatric Surgery: a Systematic Review and Meta-Analysis of the Literature. *Obes Surg.* 2016;26(12):2961-2966. doi:10.1007/s11695-016-2232-y
79. Praveenraj P, Gomes RM, Kumar S, et al. Diagnostic yield and clinical implications of preoperative upper gastrointestinal endoscopy in morbidly obese patients undergoing bariatric surgery. *J Laparoendosc Adv Surg Tech.* 2015;25(6):465-469. doi:10.1089/lap.2015.0041
80. Peromaa-Haavisto P, Victorzon M. Is routine preoperative upper GI endoscopy needed prior to gastric bypass? *Obes Surg.* 2013;23(6):736-739. doi:10.1007/s11695-013-0956-5
81. Che F, Nguyen B, Cohen A, Nguyen NT. Prevalence of hiatal hernia in the morbidly obese. *Surg Obes Relat Dis.* 2013;9(6):920-924. doi:10.1016/j.soard.2013.04.005
82. Steinbrook R. Surgery for Severe Obesity. *N Engl J Med.* 2004;350(11):1075-1079. doi:10.1177/1559827612467420
83. Küper MA, Kratt T, Kramer KM, et al. Effort, safety, and findings of routine preoperative endoscopic evaluation of morbidly obese patients undergoing bariatric surgery. *Surg Endosc.* 2010;24(8):1996-2001. doi:10.1007/s00464-010-0893-5
84. Al-Akwa A. Prevalence of Helicobacter pylori infection in a group of morbidly obese Saudi patients undergoing bariatric surgery: A preliminary report. *Saudi J Gastroenterol.* 2010;16(4):264. doi:10.4103/1319-3767.70610
85. Carabotti M, D'Ercole C, Iossa A, Corazziari E, Silecchia G, Severi C. Helicobacter pylori infection in obesity and its clinical outcome after bariatric surgery. *World J Gastroenterol.* 2014;20(3):647-653. doi:10.3748/wjg.v20.i3.647
86. Sauerland S, Angrisani L, Belachew M, et al. Obesity surgery: Evidence-based guidelines of the European Association for Endoscopic Surgery (E.A.E.S.). *Surg Endosc Other Interv Tech.* 2005;19(2):200-221. doi:10.1007/s00464-004-9194-1

87. Goday A, Castañer O, Benaiges D, et al. Can Helicobacter pylori Eradication Treatment Modify the Metabolic Response to Bariatric Surgery? *Obes Surg*. 2018;28(8):2386-2395. doi:10.1007/s11695-018-3170-7
88. U.S. Preventive Services Task Force. Recommendations for primary care practice. Recommendations. <https://www.uspreventiveservicestaskforce.org/Page/Name/recommendations>. Published 2018.
89. Himbert C, Delphan M, Scherer D, Bowers LW, Hursting S, Ulrich CM. Signals from the adipose microenvironment and the obesity-cancer link-a systematic review. *Cancer Prev Res*. 2017;10(9):494-506. doi:10.1158/1940-6207.CAPR-16-0322
90. Lashinger LM, Ford NA, Hursting SD. Interacting Inflammatory and Growth Factor Signals Underlie the Obesity-Cancer Link. *J Nutr*. 2014;144(2):109-113. doi:10.3945/jn.113.178533
91. Omalu BI, Ives DG, Buhari AM, et al. Death rates and causes of death after bariatric surgery for Pennsylvania residents, 1995 to 2004. *Arch Surg*. 2007;142(10):923-928. doi:10.1001/archsurg.142.10.929
92. Kaw R, Pasupuleti V, Wayne Overby D, et al. Inferior vena cava filters and postoperative outcomes in patients undergoing bariatric surgery: A meta-analysis. *Surg Obes Relat Dis*. 2014;10(4):725-733. doi:10.1016/j.soard.2014.04.008
93. Abou Hussein BM, Khammas A, Makki M, et al. Role of Routine Abdominal Ultrasound Before Bariatric Surgery: Review of 937 Patients. *Obes Surg*. 2018;28(9). doi:10.1007/s11695-018-3210-3
94. Mechanick JI, Kushner RF, Sugerman HJ, et al. American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery Medical guidelines for clinical practice for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric. *Obesity*. 2009;14(S1):1-83. doi:10.1038/oby.2009.28
95. Martin KA, Lee CR, Farrell TM, Moll S. Oral Anticoagulant Use After Bariatric Surgery: A Literature Review and Clinical Guidance. *Am J Med*. 2017;130(5):517-524. doi:10.1016/j.amjmed.2016.12.033
96. Geraldo M de SP, Fonseca FLA, Veiga Gouveia MR de F, Feder D. The use of drugs in patients who have undergone bariatric surgery. *Int J Gen Med*. 2014;7:219-224. doi:10.2147/IJGM.S55332
97. Padwal R, Brocks D, Sharma AM. A systematic review of drug absorption following bariatric surgery and its theoretical implications. *Obes Rev*. 2010;11(1):41-50. doi:10.1111/j.1467-789X.2009.00614.x
98. Miller AD, Smith KM. Medication and nutrient administration considerations after bariatric surgery. *Am J Heal Pharm*. 2006;63(19):1852-1857. doi:10.2146/ajhp060033
99. Estrada CA, Young JA, Nifong LW, Chitwood WR. Outcomes and perioperative hyperglycemia in patients with or without diabetes mellitus undergoing coronary artery bypass grafting. *Ann Thorac Surg*. 2003;75(5):1392-1399. doi:10.1016/S0003-4975(02)04997-4
100. Marchant MH, Viens NA, Cook C, Vail TP, Bolognesi MP. The impact of glycemic control and diabetes mellitus on perioperative outcomes after total joint arthroplasty. *J Bone Jt Surg - Ser A*. 2009;91(7):1621-1629. doi:10.2106/JBJS.H.00116
101. Walid MS, Newman BF, Yelverton JC, Nutter JP, Ajjan M, Robinson JS. Prevalence of previously unknown elevation of glycosylated hemoglobin in spine surgery patients and impact on length of stay and total cost. *J Hosp Med*. 2010;5(1):10-14. doi:10.1002/jhm.541
102. O'Sullivan CJ, Hynes N, Mahendran B, et al. Haemoglobin A1c (HbA1C) in Non-diabetic and Diabetic Vascular Patients. Is HbA1C an Independent Risk Factor and Predictor of Adverse Outcome? *Eur J Vasc Endovasc Surg*. 2006;32(2):188-197. doi:10.1016/j.ejvs.2006.01.011
103. Gustafsson UO, Thorell A, Soop M, Ljungqvist O, Nygren J. Haemoglobin A1c as a predictor of postoperative hyperglycaemia and complications after major colorectal surgery. *Br J Surg*. 2009;96(11):1358-1364. doi:10.1002/bjs.6724
104. Perna M, Romagnuolo J, Morgan K, Byrne TK, Baker M. Preoperative hemoglobin A1c and postoperative glucose control in outcomes after gastric bypass for obesity. *Surg Obes Relat Dis*. 2012;8(6):685-690. doi:10.1016/j.soard.2011.08.002
105. Chuah LL, Miras AD, Papamargaritis D, Jackson SN, Olbers T, Le Roux CW. Impact of perioperative management of glycemia in severely obese diabetic patients undergoing gastric bypass surgery. *Surg Obes Relat Dis*. 2015;11(3):578-584. doi:10.1016/j.soard.2014.11.004
106. Kahn BB, Flier JS. Obesity and insulin resistance. *J Clin Invest*. 2000;106(4):473-481. doi:10.1172/JCI10842
107. Mazer CD, Arnaout A, Connelly KA, et al. Sodium-glucose cotransporter 2 inhibitors and type 2 diabetes: clinical pearls for in-hospital initiation, in-hospital management, and postdischarge. *Curr Opin Cardiol*. 2019. doi:10.1097/HCO.0000000000000704
108. Sievenpiper JL, Chan CB, Dworatzek PD, Med CF, Williams Med SL. 2018 Clinical Practice Guidelines Nutrition Therapy Diabetes Canada Clinical Practice Guidelines Expert Committee. *Can J Diabetes*. 2018;42:564-579. doi:10.1016/j.jcjd.2017.10.009
109. Shiau JY, So DYF, Dent RR. Effects on Diabetes Medications, Weight and Glycated Hemoglobin Among Adult Patients With Obesity and Type 2 Diabetes: 6-Month Observations From a Full Meal Replacement, Low-Calorie Diet Weight Management Program. *Can J Diabetes*. 2018;42(1):56-60. doi:10.1016/j.jcjd.2017.03.006
110. Nestle Health Science. Is Optifast VLCD suitable for diabetes? <https://www.optifast-me.com/hcp/diabetes>.
111. Saiki A, Nagayama D, Ohhira M, et al. Effect of weight loss using formula diet on renal function in obese patients with diabetic nephropathy. *Int J Obes*. 2005;29(9):1115-1120. doi:10.1038/sj.ijo.0803009
112. Valenta LJ, Elias AN. Modified fasting in treatment of obesity: effects on serum lipids, electrolytes, liver enzymes, and blood pressure. *Postgrad Med*. 1986;79(4):263-267. doi:10.1080/00325481.1986.11699329
113. Couris R, Tataronis G, McCloskey W, et al. Dietary vitamin K variability affects International Normalized Ratio (INR) coagulation indices. *Int J Vitam Nutr Res*. 2006;76(2):65-74. doi:10.1024/0300-9831.76.2.65
114. Pedersen FM, Hamberg O, Hess K, Ovesen L. The effect of dietary vitamin K on warfarin-induced anticoagulation. *J Intern Med*. 1991;229(6):517-520. doi:10.1111/j.1365-2796.1991.tb00388.x